



planetmath.org

Math for the people, by the people.

Grothendieck category theorems

Canonical name	GrothendieckCategoryTheorems1
Date of creation	2013-03-22 19:22:16
Last modified on	2013-03-22 19:22:16
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	4
Author	bci1 (20947)
Entry type	Theorem
Classification	msc 18-00

# 1 Two New Theorems for Grothendieck categories

## 1.1 Introduction

The theory has its origin in the work of Grothendieck [?] who introduced the following notation of properties of abelian categories:

Ab3. An abelian category with coproducts or equivalently, a cocomplete abelian category.

Ab5. Ab3-category, in which for any directed family  $\{A_i\}_{i \in I}$  of subobjects of an arbitrary object  $X$  and for any subobject  $B$  of  $X$  the following relation holds:

$$\left(\sum_{i \in I} A_i\right) \cap B = \sum_{i \in I} (A_i \cap B)$$

Ab5-categories possessing a family of generators are called *Grothendieck categories*. They constitute a natural extension of the class of module categories, with which they share a great number of important properties.

The **Popescu-Gabriel Theorem** is generalized as follows.

**Theorem**[Popescu and Gabriel] *Let  $\mathbf{G}$  be a Grothendieck category with a family of generators  $\{U_i\}_{i \in I}$  and  $T = (-, ?) : \mathbf{G} \rightarrow \mathbf{mcAb}$  be the representation functor that takes each  $X \in \mathbf{G}$  to  $(-, X)$ , where  $\mathbf{mcAb} = \{h_{U_i} = (-, U_i)\}_{i \in I}$ . Then:*

(1.)  *$T$  is full and faithful.*

(2.)  *$T$  induces an equivalence between  $\mathbf{G}$  and the quotient category  $\mathbf{mcAbS}$ , where  $\mathbf{S}$  denotes the largest localizing subcategory in  $\mathbf{mcAb}$  for which all modules  $TX = (-, X)$  are  $\mathbf{S}$ -closed.*

This extension of the **Popescu-Gabriel Theorem** is due to *Grigory Garkusha* from the Saint-Petersburg State University, Higher Algebra and Number Theory Department, School of Mathematics and Mechanics, Bibliotechnaya Sq. 2, 198904 (Russia).

The advantage of this Theorem is that we can freely choose a family of generators  $U$  of  $\mathbf{G}$ . To be precise, if  $M$  is an arbitrary family of objects of  $\mathbf{G}$ , then the family:  $\{U_i\}_{i \in I} = U \cup M$  is also a family of generators.

We say that an object  $C$  of  $\mathbf{G}$  is  *$U$ -finitely generated* (or respectively  *$U$ -finitely presented*) if there is an epimorphism  $\eta : \psi_{i=1}^n U_i \rightarrow C$  (if there is an exact sequence  $\psi_{i=1}^n U_i \rightarrow \psi_{j=1}^m U_j \rightarrow C$ ) where  $U_i \in U$ . The full subcategory

of  $U$ -finitely generated ( $U$ -finitely presented) objects of  $\mathbf{G}$  is denoted by  $\mathbf{fg}_U \mathbf{G}$  ( $\mathbf{fp}_U \mathbf{G}$ ). When every  $U_i \in U$  is finitely generated (finitely presented), that is the functor  $(U_i, -)$  preserves direct unions (limits), we write  $\mathbf{fg}_\mathbf{G} = \mathbf{fg} \in \mathbf{G}$  ( $\mathbf{fp}_{U(\mathbf{G})} = \mathbf{fp} \in \mathbf{G}$ ). Then every Grothendieck category is locally  $U$ -finitely generated (locally  $U$ -finitely presented) which means that every object  $C$  of  $\mathbf{G}$  is a direct union (limit)

$$C = \sum_{i \in I} C_i$$

,

$$(C = \mathbf{lp}_{i \in I} C_i)$$

of  $U$ -finitely generated ( $U$ -finitely presented) objects  $C_i$ .

Recall also that a localizing subcategory  $\mathbf{S}$  of  $\mathbf{G}$  is of prefinite (finite) type provided that the inclusion functor  $\mathbf{J} : \mathbf{S} \rightarrow \mathbf{G}$  commutes with direct unions (limits). So the following proposition holds.

**Theorem**[Breitsprecher] Let  $\mathbf{G}$  be a Grothendieck category with a family of generators  $U = \{U_i\}_{i \in I}$ . Then the representation functor

$$T = (-, ?) : \mathbf{G} \rightarrow \mathbf{fp}_U \mathbf{G}^{(\text{op}, \text{Ab})}$$

defines an equivalence between  $\mathbf{G}$  and  $(\mathbf{fp}_U \mathbf{G})^{\text{op}}, \text{Ab})/\mathbf{S}$ , where  $\mathbf{S}$  is some localizing subcategory of  $\mathbf{fp}_U \mathbf{G}^{(\text{op}, \text{Ab})}$ .

Moreover,  $\mathbf{S}$  is of finite type if and only if  $\mathbf{fp}_U \mathbf{G} = \mathbf{fp} \mathbf{G}$ . In this case,  $\mathbf{G}$  is equivalent to the category

$$\mathbf{Lex}((\mathbf{fp}_U \mathbf{G})^{(\text{op}, \text{Ab})}) \text{ of contravariant left exact functors from } \mathbf{fp}_U \mathbf{G} \text{ to } \text{Ab}.$$

## References

- [Aus1] M. AUSLANDER, ‘Coherent functors’, in *Proc. Conf. on Categorical Algebra* (La Jolla, 1965), Springer, 1966, 189-231.
- [Aus2] M. AUSLANDER, ‘A functorial approach to representation theory’, *Lect. Notes Math.* 944 (1982), 105-179.
- [Aus3] M. AUSLANDER, ‘Isolated singularities and almost split sequences’, *Lect. Notes Math.* 1178 (1986), 194-242.
- [Br] S. BREITSPRECHER, ‘Lokal endlich präsentierbare Grothendieck-Kategorien’, *Mitt. Math. Sem. Giessen* 85 (1970), 1-25.

- [BD] I. BUCUR, A. DELEANU, *Introduction to the theory of categories and functors*, Wiley, London, 1968.
- [Bur] K. BURKE, ‘Some Model-Theoretic Properties of Functor Categories for Modules’, *Doctoral Thesis, University of Manchester*, 1994.
- [Fa] C. FAITH, *Algebra: rings, modules and categories*, vol. 1, Mir, Moscow, 1977 (in Russian).
- [Fr] P. FREYD, *Abelian categories*, Harper and Row, New-York, 1964.
- [Gbl] P. GABRIEL, ‘Des catégories abéliennes’, *Bull. Soc. Math. France* 90 (1962), 323-448.
- [GG1] G. A. GARKUSHA, A. I. GENERALOV, “Grothendieck categories as quotient categories of  $R - \text{mod}, Ab$ ” *Fund. Prikl. Mat.*, to appear (in Russian).
- [GG2] G. A. GARKUSHA, A. I. GENERALOV, ‘Duality for categories of finitely presented modules’, *Algebra i Analiz* to appear (in Russian).
- [Grk] A. GROTHENDIECK, ‘Sur quelques points d’algèbre homologique’, *Tohoku Math. J.* 9 (1957), 119-221.
- [GJ] L. GRUSON, C.U. JENSEN, ‘Dimensions cohomologiques reliées aux foncteurs  $lo^{(i)}$ ’, *Lect. Notes Math.* 867 (1981), 234-294.
- [Hrz] I. HERZOG, ‘The Ziegler spectrum of a locally coherent Grothendieck category’, *Proc. London Math. Soc.* 74 (1997), 503-558.
- [JL] C.U. JENSEN, M. LENZING, *Model theoretic algebra*, Logic and its Applications 2, Gordon and Breach, New York, 1989.
- [Kap] I. KAPLANSKY, *Infinite abelian groups*, Ann Arbor, University of Michigan Press, 1969.
- [Kr1] H. KRAUSE, ‘The spectrum of a locally coherent category’, *J. Pure Appl. Algebra* 114 (1997), 259-271.
- [Kr2] H. KRAUSE, ‘The spectrum of a module category’, *Habilitationsschrift, Universität Bielefeld*, 1998.

- [Kr3] H. KRAUSE, ‘Functors on locally finitely presented additive categories’, *Colloq. Math.* 75 (1998), 105-132.
- [Laz] D. LAZARD, ‘Autour de la platitude’, *Bull. Soc. Math. France* 97 (1969), 81-128.
- [PG] N. POPESCU, P. GABRIEL, ‘Caractérisation des catégories abéliennes avec générateurs et limites inductives exactes’, *C. R. Acad. Sc. Paris* 258 (1964), 4188-4190.
- [Pop] N. POPESCU, *Abelian categories with applications to rings and modules*, Academic Press, London and New-York, 1973.
- [Pr1] M. PREST, ‘Elementary torsion theories and locally finitely presented Abelian categories’, *J. Pure Appl. Algebra* 18 (1980), 205-212.
- [Pr2] M. PREST, ‘Epimorphisms of rings, interpretations of modules and strictly wild algebras’, *Comm. Algebra* 24 (1996), 517-531.
- [Pr3] M. PREST, ‘The Zariski spectrum of the category of finitely presented modules’, *preprint, University of Manchester*, 1998.
- [PRZ] M. PREST, PH. ROTHMALER, M. ZIEGLER, ‘Absolutely pure and flat modules and “indiscrete” rings’, *J. Algebra* 174 (1995), 349-372.
- [Rs] J.-E. ROOS, ‘Locally noetherian categories’, *Lect. Notes Math.* 92 (1969), 197-277.
- [Stm] B. STENSTRÖM, *Rings of quotients*, Springer-Verlag, New York and Heidelberg, 1975.
- [Zgr] M. ZIEGLER, ‘Model theory of modules’, *Ann. Pure Appl. Logic* 26 (1984), 149-213.

**more to come...**