



**planetmath.org**

Math for the people, by the people.

**resolution of a sheaf**

Canonical name	ResolutionOfASheaf
Date of creation	2013-03-22 14:42:49
Last modified on	2013-03-22 14:42:49
Owner	Simone (5904)
Last modified by	Simone (5904)
Numerical id	8
Author	Simone (5904)
Entry type	Definition
Classification	msc 18G60
Related topic	DeRhamWeilTheorem

Given a sheaf  $\mathcal{F}$  on a topological space  $X$  and a pair  $(\mathcal{F}^\bullet, d^\bullet)$  where  $\mathcal{F}^\bullet = \{\mathcal{F}^q\}_{q \in \mathbb{N}}$  is a family of sheaves on  $X$  and  $d^\bullet = \{d^q\}_{q \in \mathbb{N}}$  is a family of sheaf morphisms

$$d^q: \mathcal{F}^q \rightarrow \mathcal{F}^{q+1},$$

we say that  $(\mathcal{F}^\bullet, d^\bullet)$  is a *resolution* of  $\mathcal{F}$  if there exists an injection  $j: \mathcal{F} \rightarrow \mathcal{F}^0$  such that the sequence

$$0 \rightarrow \mathcal{F} \xrightarrow{j} \mathcal{F}^0 \xrightarrow{d^0} \mathcal{F}^1 \xrightarrow{d^1} \dots \rightarrow \mathcal{F}^q \rightarrow \dots$$

is exact.