



planetmath.org

Math for the people, by the people.

functor

Canonical name	Functor
Date of creation	2013-03-22 12:02:50
Last modified on	2013-03-22 12:02:50
Owner	nerdy2 (62)
Last modified by	nerdy2 (62)
Numerical id	7
Author	nerdy2 (62)
Entry type	Definition
Classification	msc 18-00
Synonym	covariant functor
Synonym	contravariant functor
Related topic	Endofunctor
Related topic	Monad

Given two categories \mathcal{C} and \mathcal{D} , a covariant *functor* $T : \mathcal{C} \rightarrow \mathcal{D}$ consists of an assignment for each object X of \mathcal{C} an object $T(X)$ of \mathcal{D} (i.e. a “function” $T : \text{Ob}(\mathcal{C}) \rightarrow \text{Ob}(\mathcal{D})$) together with an assignment for every morphism $f \in \text{Hom}_{\mathcal{C}}(A, B)$, to a morphism $T(f) \in \text{Hom}_{\mathcal{D}}(T(A), T(B))$, such that:

- $T(1_A) = 1_{T(A)}$ where 1_X denotes the identity morphism on the object X (in the respective category).
- $T(g \circ f) = T(g) \circ T(f)$, whenever the composition $g \circ f$ is defined.

A contravariant functor $T : \mathcal{C} \rightarrow \mathcal{D}$ is just a covariant functor $T : \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$ from the opposite category. In other words, the assignment reverses the direction of maps. If $f \in \text{Hom}_{\mathcal{C}}(A, B)$, then $T(f) \in \text{Hom}_{\mathcal{D}}(T(B), T(A))$ and $T(g \circ f) = T(f) \circ T(g)$ whenever the composition is defined (the domain of g is the same as the codomain of f).

Given a category \mathcal{C} and an object X we always have the functor $T : \mathcal{C} \rightarrow \mathbf{Sets}$ to the category of sets defined on objects by $T(A) = \text{Hom}(X, A)$. If $f : A \rightarrow B$ is a morphism of \mathcal{C} , then we define $T(f) : \text{Hom}(X, A) \rightarrow \text{Hom}(X, B)$ by $g \mapsto f \circ g$. This is a covariant functor, denoted by $\text{Hom}(X, -)$.

Similarly, one can define a contravariant functor $\text{Hom}(-, X) : \mathcal{C} \rightarrow \mathbf{Sets}$.