



planetmath.org

Math for the people, by the people.

subcategory

Canonical name	Subcategory
Date of creation	2013-03-22 14:21:58
Last modified on	2013-03-22 14:21:58
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	13
Author	CWoo (3771)
Entry type	Definition
Classification	msc 18A05
Related topic	Category
Related topic	Functor
Related topic	FaithfulFunctor
Defines	full subcategory
Defines	inclusion functor

Let \mathcal{C} be a (small) category. If \mathcal{S} is a collection of both a subset, call it $\text{Ob}(\mathcal{S})$, of objects of \mathcal{C} and a subset, call it $\text{Mor}(\mathcal{S})$, of morphisms of \mathcal{C} such that

1. For each $S \in \text{Ob}(\mathcal{S})$, the identity morphism of S , $\text{id}_S \in \text{Mor}(\mathcal{S})$;
2. For each $f \in \text{Mor}(\mathcal{S})$, $\text{domain}(f)$ and $\text{codomain}(f) \in \text{Ob}(\mathcal{S})$;
3. For every pair $f, g \in \text{Mor}(\mathcal{S})$ such that $f \circ g$ exists, then $f \circ g \in \text{Mor}(\mathcal{S})$.

Then \mathcal{S} is readily seen to be a category. It is called a *subcategory* of the category \mathcal{C} .

Given a category \mathcal{C} and a subcategory \mathcal{S} of \mathcal{C} , a map

$$\text{Incl} : \mathcal{S} \hookrightarrow \mathcal{C}$$

that sends each object of \mathcal{S} to itself (in \mathcal{C}), and each morphism of \mathcal{S} to itself (in \mathcal{C}), is a functor. Incl is called the *inclusion functor*, or an *embedding*. This inclusion functor is a faithful functor. If it is also <http://planetmath.org/fullfunctorfull>, then we call the corresponding subcategory \mathcal{S} a *full subcategory* of \mathcal{C} . In other words, if \mathcal{S} is a full subcategory of \mathcal{C} , then

$$\text{hom}_{\mathcal{C}}(S_1, S_2) = \text{hom}_{\mathcal{S}}(S_1, S_2)$$

for pair of $S_1, S_2 \in \text{Ob}(\mathcal{S})$.

Remarks

1. Let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a full and faithful functor. Then $T(\mathcal{C})$ is a full subcategory of \mathcal{D} .
2. Again, let $T : \mathcal{C} \rightarrow \mathcal{D}$ be a full and faithful functor. If \mathcal{S} is a full subcategory of \mathcal{D} , then $T^{-1}(\mathcal{S})$ defined by:

- $\text{Ob}(T^{-1}(\mathcal{S})) := \{C \in \text{Ob}(\mathcal{C}) \mid T(C) \in \text{Ob}(\mathcal{S})\}$
- $\text{Mor}(T^{-1}(\mathcal{S})) := \{f \in \text{Mor}(\mathcal{C}) \mid T(f) \in \text{Mor}(\mathcal{S})\}$

is a subcategory of \mathcal{C} .

Examples of Subcategories

1. In **Set**, the category of finite sets is a full subcategory, and so is the category of k -element sets, where k is any (possibly infinite) cardinality. If k is finite, then every morphism in the subcategory is invertible.

2. In **Top**, we have the full subcategories whose objects are Euclidean spaces, compact spaces, or Hausdorff spaces.
3. In **Grp**, there is the full subcategory whose objects are abelian groups with additive homomorphisms.
4. **Grp** is in fact a subcategory of the category of topological groups, since every group may be viewed as a topological group with the discrete topology.
5. In , there are the subcategories of commutative rings, matrix rings, or fields. Note that **Field** is not a full subcategory of **Ring**, since the ring homomorphism that maps every element to 0 is not a field homomorphism.

References

- [1] S. Mac Lane, *Categories for the Working Mathematician* (2nd edition), Springer-Verlag, 1997.