



planetmath.org

Math for the people, by the people.

equivalence of categories

Canonical name	EquivalenceOfCategories
Date of creation	2013-03-22 13:29:34
Last modified on	2013-03-22 13:29:34
Owner	mhale (572)
Last modified by	mhale (572)
Numerical id	7
Author	mhale (572)
Entry type	Definition
Classification	msc 18A40
Related topic	EssentiallySurjective

Let C and D be two categories with functors $F: C \rightarrow D$ and $G: D \rightarrow C$. The functors F and G are an **equivalence of categories** if there are natural isomorphisms $FG \cong \text{id}_D$ and $GF \cong \text{id}_C$.

Note, F is left adjoint to G , and G is right adjoint to F as

$$\text{hom}_D(F(c), d) \xrightarrow{G} \text{hom}_C(GF(c), G(d)) \longleftrightarrow \text{hom}_C(c, G(d)).$$

And, F is right adjoint to G , and G is left adjoint to F as

$$\text{hom}_C(G(d), c) \xrightarrow{F} \text{hom}_D(FG(d), F(c)) \longleftrightarrow \text{hom}_D(d, F(c)).$$

In practical terms, two categories are equivalent if there is a <http://planetmath.org/FullFunctor> faithful functor $F: C \rightarrow D$, such that every object $d \in D$ is isomorphic to an object $F(c)$, for some $c \in C$.