



planetmath.org

Math for the people, by the people.

product of categories

Canonical name	ProductOfCategories
Date of creation	2013-03-22 16:37:05
Last modified on	2013-03-22 16:37:05
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	10
Author	CWoo (3771)
Entry type	Definition
Classification	msc 18A05
Related topic	CategoricalDirectProduct
Related topic	DisjointUnionOfCategories
Defines	product category
Defines	projection functor

There are occasions when we need to consider objects from different categories being paired up. For example, if  $\mathcal{C}$  is a category, then  $\text{hom}(A, B)$  where  $A, B$  are objects of  $\mathcal{C}$  is a set, and we can think of  $\text{hom}$  as a functor from  $\mathcal{C}^{\text{op}} \times \mathcal{C}$  to the category of sets. But what is  $\mathcal{C}^{\text{op}} \times \mathcal{C}$  exactly? We will give this a formal definition presently.

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories. Define the *Cartesian product*  $\mathcal{C} \times \mathcal{D}$  of  $\mathcal{C}$  and  $\mathcal{D}$  as the following pair  $(O, M)$ , where

- $O$  is the class consisting of ordered pairs  $(X, Y)$ , where  $X$  is an object in  $\mathcal{C}$  and  $Y$  is an object in  $\mathcal{D}$
- $M$  is the class consisting of ordered pairs  $(f, g)$ , where  $f$  is a morphism in  $\mathcal{C}$  and  $g$  is a morphism in  $\mathcal{D}$ .

There is a category structure on  $\mathcal{C} \times \mathcal{D}$ . But several things need to be defined first.

1. Elements of  $O$  are called the *objects* of  $\mathcal{C} \times \mathcal{D}$  and elements of  $M$  are called the *morphisms* of  $\mathcal{C} \times \mathcal{D}$ . For each morphism  $(f, g) \in M$ , we define the domain and codomain operations

$$\text{dom}(f, g) := (\text{dom}(f), \text{dom}(g)) \quad \text{and} \quad \text{cod}(f, g) := (\text{cod}(f), \text{cod}(g)).$$

Note that for simplicity, we have used the same symbol  $\text{dom}$  and  $\text{cod}$  for  $\mathcal{C}$ ,  $\mathcal{D}$ , and  $\mathcal{C} \times \mathcal{D}$ .

2. Next, for each pair of objects  $A, B \in \mathcal{C} \times \mathcal{D}$ , we have a set  $\text{hom}(A, B)$  consisting of all morphisms in  $\mathcal{C} \times \mathcal{D}$  whose domain is  $A$  and codomain is  $B$ . Note that  $\text{hom}(A, B)$  is a set because it is  $\text{hom}(X, Z) \times \text{hom}(Y, T)$ , where  $A = (X, Y)$  and  $B = (Z, T)$  and each component in the product is assumed to be a set.
3. Finally, for objects  $A, B, C$  in  $\mathcal{C} \times \mathcal{D}$ , we have a function  $\circ$  called composition:

$$\circ : \text{hom}(A, B) \times \text{hom}(B, C) \rightarrow \text{hom}(A, C).$$

To define  $\circ$ , write each object  $A, B, C$  as ordered pairs:  $A = (X, Y)$ ,  $B = (Z, T)$ ,  $C = (U, V)$ . In addition, let  $\alpha = (f, g) \in \text{hom}(A, B)$  and  $\beta = (p, q) \in \text{hom}(B, C)$ . Then

$$\circ(\alpha, \beta) := (\circ_1(f, p), \circ_2(g, q)),$$

where  $\circ_1$  and  $\circ_2$  are compositions defined in  $\mathcal{C}$  and  $\mathcal{D}$  respectively, such that

$$\circ_1 : \text{hom}(X, Z) \times \text{hom}(Z, U) \rightarrow \text{hom}(X, U) \text{ and } \circ_2 : \text{hom}(Y, T) \times \text{hom}(T, V) \rightarrow \text{hom}(Y, V).$$

As usual, we write  $\beta \circ \alpha$  for  $\circ(\alpha, \beta)$ .

4. Now, it is not hard to see that  $\mathcal{C} \times \mathcal{D}$  with  $\circ$  is a category. For example, let us verify that  $(A, B) \neq (C, D)$  implies  $\text{hom}(A, B) \cap \text{hom}(C, D) = \emptyset$ . Write  $A = (X, Y)$ ,  $B = (Z, S)$ ,  $C = (T, U)$  and  $D = (V, W)$ . Suppose  $\alpha = (f, g) \in \text{hom}(A, B) \cap \text{hom}(C, D)$ . Then  $f \in \text{hom}(X, Z) \cap \text{hom}(T, V)$  and  $g \in \text{hom}(Y, S) \cap \text{hom}(U, W)$ . But this implies  $X = T$ ,  $Z = V$ ,  $Y = U$ , and  $S = W$ . So  $A = (X, Y) = (T, U) = C$  and  $B = (Z, S) = (V, W) = D$ .

#### Remarks.

- The above construction can be generalized to  $n$ -fold Cartesian products. If  $\mathcal{C}_1, \dots, \mathcal{C}_n$  be categories. Then  $\mathcal{C} := \mathcal{C}_1 \times \dots \times \mathcal{C}_n$  can be defined much the same way as in the case  $n = 2$ .  $\mathcal{C}$  is a category and is sometimes written  $\prod_{i=1}^n \mathcal{C}_i$ .
- Associated with this product, we can form  $n$  (covariant) functors called *projection functors*  $\Pi_i : \mathcal{C} \rightarrow \mathcal{C}_i$ , given by  $\Pi_i(A) = A_i$  and  $\Pi_i(\alpha) = \alpha_i$ , where  $A = (A_1, \dots, A_n)$  and  $\alpha = (\alpha_1, \dots, \alpha_n)$ .
- The product  $\mathcal{C}$  of  $\mathcal{C}_i$  also enjoys the universal property that for every category  $\mathcal{D}$  and functors  $F_i : \mathcal{D} \rightarrow \mathcal{C}_i$ , there is a unique functor  $F : \mathcal{D} \rightarrow \mathcal{C}$  such that  $\Pi_i \circ F = F_i$  (in other words,  $F_i$  factors through  $F$ ).
- In fact, any category that enjoys the universal property described above is naturally equivalent to the product of  $\mathcal{C}_i$ . We may actually define product category this way, and then prove its existence using the construction that is given as the definition at the beginning of this article.
- More generally, we can define arbitrary (direct) product of categories. The definition is completely similar to the one above. If  $\{\mathcal{C}_i \mid i \in I\}$  is a family of categories indexed by a set  $I$ , we often write  $\prod_{i \in I} \mathcal{C}_i$  as the product category. Objects and morphisms are written  $(A_i)_{i \in I}$  and  $(\alpha_i)_{i \in I}$  respectively. When all the  $\mathcal{C}_i$  are identical, say, equal to  $\mathcal{C}$ , we also write the product as  $\mathcal{C}^I$ , and call it the *I-fold direct product of  $\mathcal{C}$* .

- The existence of the product of categories indexed by an arbitrary set shows that the category of (small) categories **Cat** has products.
- Let  $\mathcal{C} = \mathcal{D} \times \mathcal{E}$ . Then we may identify  $\mathcal{D}$  as a subcategory of  $\mathcal{C}$ : for each object  $E$  in  $\mathcal{E}$ , define  $F_E : \mathcal{D} \rightarrow \mathcal{C}$ , by  $F_E(A) := (A, E)$  and  $F(\alpha) = (\alpha, 1_E)$ . Then  $F_E$  is a faithful functor. The image  $\mathcal{C}_E$  of  $F_E$  (with objects  $(A, E)$  and morphisms  $(\alpha, 1_E)$ ) is a subcategory of  $\mathcal{C}$ . It is not hard to see that  $\mathcal{D}$  and  $\mathcal{C}_E$  are <http://planetmath.org/CategoryIsomorphism> isomorphic as categories.
- The above also shows that for any objects  $A, B$  in  $\mathcal{E}$ ,  $\mathcal{C}_A$  and  $\mathcal{C}_B$  are isomorphic.