

## planetmath.org

Math for the people, by the people.

## generator of a category

Canonical name GeneratorOfACategory
Date of creation 2013-03-22 18:21:04
Last modified on 2013-03-22 18:21:04

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 10

Author CWoo (3771) Entry type Definition Classification msc 18A99

Related topic GrothendieckCategory

Defines generator
Defines generating set
Defines progenerator

Let  $\mathcal{C}$  be a category, and  $f, g: A \to B$  a pair of distinct morphisms. A morphism  $h: X \to A$  is said to distinguish or separate f and g if  $f \circ h \neq g \circ h$ . For example, if  $f \neq g: A \to B$ , then  $1_A$  on A distinguishes f and g.

A set  $S = \{X_i \mid i \in I\}$  of objects (indexed by a set I) is called a generating set of C if any pair of distinct morphisms  $f, g : A \to B$  can be distinguished by a morphism with domain in S and codomain A. In other words, there is  $h : X_i \to A$  for some  $i \in I$ , such that  $f \circ h \neq g \circ h$ . If  $\{X\}$  is a generating family of C, then X is called a generator of C. Any set of morphisms containing a generator is a generating set.

## Examples

- 1. In **Set**, the category of sets, any singleton is a generator. Suppose  $f, g: A \to B$  are distinct functions, so that  $f(x) \neq g(x)$  for some  $x \in A$ . Let  $\{y\}$  be any singleton. Then  $h: \{y\} \to A$  defined by h(y) = x is the function distinguishing f and g: for  $f \circ h(y) = f(x) \neq g(x) = g \circ h(y)$ .
- 2. In **Rng**, the category of rings, the ring  $\mathbb{Z}$  is a generator. If  $f, g : R \to S$  are distinct ring homomorphisms, say,  $f(r) \neq g(r)$  for some  $r \in R$ . Then the ring homomorphism  $h : \mathbb{Z} \to R$  given by h(1) = r distinguishes f and g.

**Remark**. A projective object that is also a generator is called a *progenerator*.

## References

[1] F. Borceux Basic Category Theory, Handbook of Categorical Algebra I, Cambridge University Press, Cambridge (1994)