



planetmath.org

Math for the people, by the people.

algebra formed from a category

Canonical name	AlgebraFormedFromACategory
Date of creation	2013-03-22 16:30:34
Last modified on	2013-03-22 16:30:34
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	6
Author	rspuzio (6075)
Entry type	Definition
Classification	msc 18A05

Given a category \mathcal{C} and a ring R , one can construct an algebra \mathcal{A} as follows. Let \mathcal{A} be the set of all formal finite linear combinations of the form

$$\sum_i c_i e_{a_i, b_i, \mu_i},$$

where the coefficients c_i lie in R and, to every pair of objects a and b of \mathcal{C} and every morphism μ from a to b , there corresponds a basis element $e_{a, b, \mu}$. Addition and scalar multiplication are defined in the usual way. Multiplication of elements of \mathcal{A} may be defined by specifying how to multiply basis elements. If $b \neq c$, then set $e_{a, b, \phi} \cdot e_{c, d, \psi} = 0$; otherwise set $e_{a, b, \phi} \cdot e_{b, c, \psi} = e_{a, c, \psi \circ \phi}$. Because of the associativity of composition of morphisms, \mathcal{A} will be an associative algebra over R .

Two instances of this construction are worth noting. If G is a group, we may regard G as a category with one object. Then this construction gives us the group algebra of G . If P is a partially ordered set, we may view P as a category with at most one morphism between any two objects. Then this construction provides us with the incidence algebra of P .