



planetmath.org

Math for the people, by the people.

groupoid action

Canonical name	GroupoidAction
Date of creation	2013-03-22 19:19:23
Last modified on	2013-03-22 19:19:23
Owner	bci1 (20947)
Last modified by	bci1 (20947)
Numerical id	9
Author	bci1 (20947)
Entry type	Definition
Classification	msc 22A22
Classification	msc 18B40
Synonym	action
Related topic	GroupAction
Related topic	Groupoid
Related topic	GroupoidRepresentation4
Defines	anchor map

**Definition 0.1.** Let  $\mathcal{G}$  be a groupoid and  $X$  a topological space. A *groupoid action*, or  $\mathcal{G}$ -action, on  $X$  is given by two maps: the *anchor map*  $\pi : X \longrightarrow G_0$  and a map  $\mu : X \times_{G_0} G_1 \longrightarrow X$ , with the latter being defined on pairs  $(x, g)$  such that  $\pi(x) = t(g)$ , written as  $\mu(x, g) = xg$ . The two maps are subject to the following conditions:

- $\pi(xg) = s(g)$ ,
- $xu(\pi(x)) = x$ , and
- $(xg)h = x(gh)$ , whenever the operations are defined.

**Note:** The groupoid action generalizes the concept of group action in a non-trivial way.