

## planetmath.org

Math for the people, by the people.

## topological \*-algebra

Canonical name Topological algebra
Date of creation 2013-03-22 14:45:38
Last modified on 2013-03-22 14:45:38

Owner HkBst (6197) Last modified by HkBst (6197)

Numerical id 12

Author HkBst (6197)
Entry type Definition
Classification msc 22A30
Classification msc 16W80
Classification msc 16W10
Classification msc 46K05
Classification msc 46H35

Synonym topological \*-algebra

Related topic BanachAlgebra

Related topic WeakHopfCAlgebra2
Related topic VonNeumannAlgebra
Defines involution \*-algebra

Defines \*-algebra

**Definition (Involution)** An involution on an algebra A over an http://planetmath.org/Involution field F is a map  $\cdot^*: A \to A: a \mapsto a^*$  such that for every  $\{a,b\} \subset A$  and  $\lambda \in F$  we have

- 1.  $a^{**} = a$ ,
- 2.  $(ab)^* = b^*a^*$  and
- 3.  $(\lambda a + b)^* = \lambda^* a^* + b^*$ , where  $\lambda^*$  denotes the http://planetmath.org/InvolutaryRinginvoluti of  $\lambda$  in F.

**Definition** (\*-Algebra) A \*-algebra is an algebra with an involution. **Definition** (**Topological** \*-algebra) A topological \*-algebra is a \*-algebra which is also a topological vector space such that its algebra multiplication and involution are continuous.

## 0.0.1 Remarks:

- \*-algebras are a particular of involutory rings.
- The involutory field F is often taken as  $\mathbb{C}$ , where the involution is given by complex conjugation. In this case, condition 3 could be rewritten as:

3. 
$$(\lambda a + b)^* = \overline{\lambda} a^* + b^*$$

• Banach \*-algebras are topological \*-algebras.