



Math for the people, by the people.

one-parameter subgroup

| | |
|------------------|----------------------|
| Canonical name | OneparameterSubgroup |
| Date of creation | 2013-03-22 14:54:01 |
| Last modified on | 2013-03-22 14:54:01 |
| Owner | CWoo (3771) |
| Last modified by | CWoo (3771) |
| Numerical id | 7 |
| Author | CWoo (3771) |
| Entry type | Definition |
| Classification | msc 22E15 |
| Classification | msc 22E10 |
| Synonym | 1-parameter subgroup |

Let G be a Lie Group. A *one-parameter subgroup* of G is a group homomorphism

$$\phi: \mathbb{R} \rightarrow G$$

that is also a differentiable map at the same time. We view \mathbb{R} additively and G multiplicatively, so that $\phi(r + s) = \phi(r)\phi(s)$.

Examples.

1. If $G = \text{GL}(n, k)$, where $k = \mathbb{R}$ or \mathbb{C} , then any one-parameter subgroup has the form

$$\phi(t) = e^{tA},$$

where $A = \frac{d\phi}{dt}(0)$ is an $n \times n$ matrix over k . The matrix A is just a tangent vector to the Lie group $\text{GL}(n, k)$. This property establishes the fact that there is a one-to-one correspondence between one-parameter subgroups and tangent vectors of $\text{GL}(n, k)$. The same relationship holds for a general Lie group. The one-to-one correspondence between tangent vectors at the identity (the Lie algebra) and one-parameter subgroups is established via the exponential map instead of the matrix exponential.

2. If $G = \text{O}(n, \mathbb{R}) \subseteq \text{GL}(n, \mathbb{R})$, the orthogonal group over \mathbb{R} , then any one-parameter subgroup has the same form as in the example above, except that A is skew-symmetric: $A^T = -A$.
3. If $G = \text{SL}(n, \mathbb{R}) \subseteq \text{GL}(n, \mathbb{R})$, the special linear group over \mathbb{R} , then any one-parameter subgroup has the same form as in the example above, except that $\text{tr}(A) = 0$, where tr is the trace operator.
4. If $G = \text{U}(n) = \text{O}(n, \mathbb{C}) \subseteq \text{GL}(n, \mathbb{C})$, the unitary group over \mathbb{C} , then any one-parameter subgroup has the same form as in the example above, except that A is <http://planetmath.org/SkewHermitianMatrix> skew-Hermitian: $A = -A^* = -\overline{A}^T$ and $\text{tr}(A) = 0$.