



**planetmath.org**

Math for the people, by the people.

## **another proof of Jensen's inequality**

Canonical name	AnotherProofOfJensensInequality
Date of creation	2013-03-22 15:52:53
Last modified on	2013-03-22 15:52:53
Owner	Andrea Ambrosio (7332)
Last modified by	Andrea Ambrosio (7332)
Numerical id	12
Author	Andrea Ambrosio (7332)
Entry type	Proof
Classification	msc 26D15
Classification	msc 39B62

First of all, it's clear that defining

$$\lambda_k = \frac{\mu_k}{\sum_{k=1}^n \mu_k}$$

we have

$$\sum_{k=1}^n \lambda_k = 1$$

so it will be enough to prove only the simplified version.

Let's proceed by induction.

1)  $n = 2$ ; we have to show that, for any  $x_1$  and  $x_2$  in  $[a, b]$ ,

$$f(\lambda_1 x_1 + \lambda_2 x_2) \leq \lambda_1 f(x_1) + \lambda_2 f(x_2).$$

But, since  $\lambda_1 + \lambda_2$  must be equal to 1, we can put  $\lambda_2 = 1 - \lambda_1$ , so that the thesis becomes

$$f(\lambda_1 x_1 + (1 - \lambda_1) x_2) \leq \lambda_1 f(x_1) + (1 - \lambda_1) f(x_2),$$

which is true by definition of a convex function.

2) Taking as true that  $f(\sum_{k=1}^{n-1} \mu_k x_k) \leq \sum_{k=1}^{n-1} \mu_k f(x_k)$ , where  $\sum_{k=1}^{n-1} \mu_k = 1$ , we have to prove that

$$f\left(\sum_{k=1}^n \lambda_k x_k\right) \leq \sum_{k=1}^n \lambda_k f(x_k),$$

where  $\sum_{k=1}^n \lambda_k = 1$ .

First of all, let's observe that

$$\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} = \frac{(\sum_{k=1}^{n-1} \lambda_k) - \lambda_n}{1 - \lambda_n} = \frac{1 - \lambda_n}{1 - \lambda_n} = 1$$

and that if all  $x_k \in [a, b]$ ,  $\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k$  belongs to  $[a, b]$  as well. In fact,  $\frac{\lambda_k}{1 - \lambda_n}$  being non-negative,

$$a \leq x_k \leq b \Rightarrow \frac{\lambda_k}{1 - \lambda_n} a \leq \frac{\lambda_k}{1 - \lambda_n} x_k \leq \frac{\lambda_k}{1 - \lambda_n} b,$$

and, summing over  $k$ ,

$$a \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} \leq \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k \leq b \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n},$$

that is

$$a \leq \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k \leq b.$$

We have, by definition of a convex function:

$$\begin{aligned} f\left(\sum_{k=1}^n \lambda_k x_k\right) &= f\left(\sum_{k=1}^{n-1} \lambda_k x_k + \lambda_n x_n\right) \\ &= f\left((1 - \lambda_n) \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k + \lambda_n x_n\right) \\ &\leq (1 - \lambda_n) f\left(\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k\right) + \lambda_n f(x_n). \end{aligned}$$

But, by inductive hypothesis, since  $\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} = 1$ , we have:

$$f\left(\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k\right) \leq \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} f(x_k),$$

so that

$$\begin{aligned} f\left(\sum_{k=1}^n \lambda_k x_k\right) &\leq (1 - \lambda_n) f\left(\sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} x_k\right) + \lambda_n f(x_n) \\ &\leq (1 - \lambda_n) \sum_{k=1}^{n-1} \frac{\lambda_k}{1 - \lambda_n} f(x_k) + \lambda_n f(x_n) \\ &= \sum_{k=1}^{n-1} \lambda_k f(x_k) + \lambda_n f(x_n) \\ &= \sum_{k=1}^n \lambda_k f(x_k) \end{aligned}$$

which is the thesis.