



**planetmath.org**

Math for the people, by the people.

## higher order derivatives of sine and cosine

|                  |                                       |
|------------------|---------------------------------------|
| Canonical name   | HigherOrderDerivativesOfSineAndCosine |
| Date of creation | 2013-03-22 14:45:16                   |
| Last modified on | 2013-03-22 14:45:16                   |
| Owner            | pahio (2872)                          |
| Last modified by | pahio (2872)                          |
| Numerical id     | 13                                    |
| Author           | pahio (2872)                          |
| Entry type       | Derivation                            |
| Classification   | msc 26B05                             |
| Classification   | msc 46G05                             |
| Classification   | msc 26A24                             |
| Related topic    | FractionalDifferentiation             |
| Related topic    | HigherOrderDerivatives                |
| Related topic    | ExampleOfTaylorPolynomialsForSinX     |
| Related topic    | CosineAtMultiplesOfStraightAngle      |

One may consider the sine and cosine either as <http://planetmath.org/RealFunctionreal> or complex functions. In both cases they are everywhere smooth, having the derivatives of all <http://planetmath.org/OrderOfDerivativeorders> in every point. The formulae

$$\frac{d^n}{dx^n} \sin x = \sin \left( x + n \cdot \frac{\pi}{2} \right)$$

and

$$\frac{d^n}{dx^n} \cos x = \cos \left( x + n \cdot \frac{\pi}{2} \right),$$

where  $n = 0, 1, 2, \dots$  (the derivative of the 0<sup>th</sup> order means the function itself), can be proven by induction on  $n$ . Another possibility is to utilize Euler's formula, obtaining

$$\frac{d^n}{dx^n} \cos x + i \frac{d^n}{dx^n} \sin x = \frac{d^n}{dx^n} e^{ix} = e^{ix} i^n = e^{ix + in \frac{\pi}{2}} = \cos \left( x + n \cdot \frac{\pi}{2} \right) + i \sin \left( x + n \cdot \frac{\pi}{2} \right);$$

here one has to compare the <http://planetmath.org/ComplexFunctionreal> and imaginary parts – supposing that  $x$  is real.