



Math for the people, by the people.

differential equations for x^x

Canonical name	DifferentialEquationsForXx
Date of creation	2013-03-22 17:24:37
Last modified on	2013-03-22 17:24:37
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	6
Author	rspuzio (6075)
Entry type	Derivation
Classification	msc 26A99

In this entry, we will derive differential equations satisfied by the function x^x .¹ We begin by computing its derivative. To do this, we write $x^x = e^{x \log x}$ and apply the chain rule:

$$\frac{d}{dx}x^x = \frac{d}{dx}e^{x \log x} = e^{x \log x}(1 + \log x) = x^x(1 + \log x)$$

Set $y = x^x$. Then we have $y'/y = 1 + \log x$. Taking another derivative, we have

$$\frac{d}{dx} \left(\frac{y'}{y} \right) = \frac{1}{x}.$$

Applying the quotient rule and simplifying, this becomes

$$yy'' - (y')^2 - y^2/x = 0.$$

It is also possible to derive an equation in which x does not appear. We start by noting that, if $z = 1/x$, then $z' + z^2 = 0$. If, as above, $y = x^x$, we have $(d/dx)(y'/y) = z$. Combining equations,

$$\frac{d^2}{dx^2} \left(\frac{y'}{y} \right) + \left(\frac{d}{dx} \left(\frac{y'}{y} \right) \right)^2 = 0;$$

applying the quotient rule and simplifying,

$$y^3y''' - y^2(y'')^2 + 2y(y')^2y'' - 3y^2y'y'' - (y')^4 + 2y(y')^3 = 0.$$

¹In this entry, we restrict x , and hence x^x to be strictly positive real numbers, hence it is justified to divide by these quantities.