



planetmath.org

Math for the people, by the people.

logarithmically convex function

Canonical name	LogarithmicallyConvexFunction
Date of creation	2013-03-22 14:13:33
Last modified on	2013-03-22 14:13:33
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	7
Author	jirka (4157)
Entry type	Definition
Classification	msc 26A51
Synonym	logarithmically convex
Synonym	log-convex function
Synonym	log-convex
Synonym	log convex function
Synonym	log convex
Related topic	ConvexFunction
Related topic	BohrMollerupTheorem
Related topic	HadamardThreeCircleTheorem

Definition. A function $f: [a, b] \rightarrow \mathbb{R}$ such that $f(x) > 0$ for all x is said to be *logarithmically convex* if $\log f(x)$ is a convex function.

It is easy to see that a logarithmically convex function is a convex function, but the converse is not true. For example $f(x) = x^2$ is a convex function, but $\log f(x) = \log x^2 = 2 \log x$ is not a convex function and thus $f(x) = x^2$ is not logarithmically convex. On the other hand e^{x^2} is logarithmically convex since $\log e^{x^2} = x^2$ is convex. A less trivial example of a logarithmically convex function is the gamma function, if restricted to the positive reals.

The definition is easily extended to functions $f: U \subset \mathbb{R} \rightarrow \mathbb{R}$, for any connected set U (where still we have $f > 0$), in the obvious way. Such a function is logarithmically convex if it is logarithmically convex on all intervals $[a, b] \subset U$.

References

- [1] John B. Conway. . Springer-Verlag, New York, New York, 1978.