



Math for the people, by the people.

## proof of Fermat's Theorem (stationary points)

Canonical name	ProofOfFermatsTheoremstationaryPoints
Date of creation	2013-03-22 13:45:09
Last modified on	2013-03-22 13:45:09
Owner	paolini (1187)
Last modified by	paolini (1187)
Numerical id	5
Author	paolini (1187)
Entry type	Proof
Classification	msc 26A06

Suppose that  $x_0$  is a local maximum (a similar proof applies if  $x_0$  is a local minimum). Then there exists  $\delta > 0$  such that  $(x_0 - \delta, x_0 + \delta) \subset (a, b)$  and such that we have  $f(x_0) \geq f(x)$  for all  $x$  with  $|x - x_0| < \delta$ . Hence for  $h \in (0, \delta)$  we notice that it holds

$$\frac{f(x_0 + h) - f(x_0)}{h} \leq 0.$$

Since the limit of this ratio as  $h \rightarrow 0^+$  exists and is equal to  $f'(x_0)$  we conclude that  $f'(x_0) \leq 0$ . On the other hand for  $h \in (-\delta, 0)$  we notice that

$$\frac{f(x_0 + h) - f(x_0)}{h} \geq 0$$

but again the limit as  $h \rightarrow 0^+$  exists and is equal to  $f'(x_0)$  so we also have  $f'(x_0) \geq 0$ .

Hence we conclude that  $f'(x_0) = 0$ .

To prove the second part of the statement (when  $x_0$  is equal to  $a$  or  $b$ ), just notice that in such points we have only one of the two estimates written above.