



Math for the people, by the people.

## proof of mean value theorem

Canonical name	ProofOfMeanValueTheorem
Date of creation	2013-03-22 12:40:57
Last modified on	2013-03-22 12:40:57
Owner	Andrea Ambrosio (7332)
Last modified by	Andrea Ambrosio (7332)
Numerical id	5
Author	Andrea Ambrosio (7332)
Entry type	Proof
Classification	msc 26A06

Define  $h(x)$  on  $[a, b]$  by

$$h(x) = f(x) - f(a) - \left( \frac{f(b) - f(a)}{b - a} \right) (x - a)$$

Clearly,  $h$  is continuous on  $[a, b]$ , differentiable on  $(a, b)$ , and

$$\begin{aligned} h(a) &= f(a) - f(a) = 0 \\ h(b) &= f(b) - f(a) - \left( \frac{f(b) - f(a)}{b - a} \right) (b - a) = 0 \end{aligned}$$

Notice that  $h$  satisfies the conditions of Rolle's Theorem. Therefore, by Rolle's Theorem there exists  $c \in (a, b)$  such that  $h'(c) = 0$ .

However, from the definition of  $h$  we obtain by differentiation that

$$h'(x) = f'(x) - \frac{f(b) - f(a)}{b - a}$$

Since  $h'(c) = 0$ , we therefore have

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

as required.

## References

- [1] Michael Spivak, *Calculus*, 3rd ed., Publish or Perish Inc., 1994.