



Math for the people, by the people.

Clairaut's theorem

Canonical name	ClairautsTheorem
Date of creation	2013-03-22 13:53:44
Last modified on	2013-03-22 13:53:44
Owner	Mathprof (13753)
Last modified by	Mathprof (13753)
Numerical id	18
Author	Mathprof (13753)
Entry type	Theorem
Classification	msc 26B12
Synonym	equality of mixed partials

Clairaut's Theorem. *If $\mathbf{f}: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a function whose second partial derivatives exist and are continuous on a set $S \subseteq \mathbb{R}^n$, then*

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}$$

on S , where $1 \leq i, j \leq n$.

This theorem is commonly referred to as *the equality of mixed partials*. It is usually first presented in a vector calculus course, and is useful in this context for proving basic properties of the interrelations of gradient, divergence, and curl. For example, if $\mathbf{F}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a function satisfying the hypothesis, then $\nabla \cdot (\nabla \times \mathbf{F}) = 0$. Or, if $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function satisfying the hypothesis, then $\nabla \times \nabla f = \mathbf{0}$.