



**planetmath.org**

Math for the people, by the people.

**ruler function**

Canonical name	RulerFunction
Date of creation	2013-03-22 18:23:55
Last modified on	2013-03-22 18:23:55
Owner	yesitis (13730)
Last modified by	yesitis (13730)
Numerical id	5
Author	yesitis (13730)
Entry type	Definition
Classification	msc 26A99

The ruler function  $f$  on the real line is defined as follows:

$$f(x) = \begin{cases} 0, & x \text{ is irrational;} \\ 1/n, & x = m/n, m \text{ and } n \text{ are relatively primes.} \end{cases} \quad (1)$$

Given a rational number  $\frac{m}{n}$  in lowest terms,  $n$  positive, the ruler function outputs the size (length) of a piece resulting from equally subdividing the unit interval into  $n$ , the number in the denominator, parts. It “ignores” inputs of irrational functions, sending them to 0.

The ruler function is so termed because it resembles a ruler. The following picture might be helpful: if  $\frac{m}{n}$  in lowest terms is a reasonably small rational number (which we assume positive). Then it can be “read off” on a ruler whose intervals of one unit size are each equally subdivided into  $n$  parts measuring  $\frac{1}{n}$  units each by

1. running one’s finger through until the integer preceding it and then
2. running through to the subsequent  $r$ th subunit, “left-over” from the division of  $m$  by  $n$ .

On the other hand, an irrational number can not be read off from any ruler no matter how fine we subdivide a unit interval in any ruler.

## References

- [1] Dunham, W., *Nondifferentiability of the Ruler Function*, Mathematics Magazine, Mathematical Association of America, 2003.
- [2] Heuer, G.A., *Functions Continuous at the Irrationals and Discontinuous at the Rationals*, The American Mathematical Monthly, Mathematical Association of America, 1965.