



planetmath.org

Math for the people, by the people.

monotone convergence theorem

Canonical name	MonotoneConvergenceTheorem
Date of creation	2013-03-22 12:47:27
Last modified on	2013-03-22 12:47:27
Owner	Koro (127)
Last modified by	Koro (127)
Numerical id	9
Author	Koro (127)
Entry type	Theorem
Classification	msc 26A42
Classification	msc 28A20
Synonym	Lebesgue's monotone convergence theorem
Synonym	Beppo Levi's theorem
Related topic	DominatedConvergenceTheorem
Related topic	FatousLemma

Let  $X$  be a measure space, and let  $0 \leq f_1 \leq f_2 \leq \cdots$  be a monotone increasing sequence of nonnegative measurable functions. Let  $f: X \rightarrow \mathbb{R} \cup \{\infty\}$  be the function defined by  $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ . Then  $f$  is measurable, and

$$\lim_{n \rightarrow \infty} \int_X f_n = \int_X f.$$

**Remark.** This theorem is the first of several theorems which allow us to “exchange integration and limits”. It requires the use of the Lebesgue integral: with the Riemann integral, we cannot even formulate the theorem, lacking, as we do, the concept of “almost everywhere”. For instance, the characteristic function of the rational numbers in  $[0, 1]$  is not Riemann integrable, despite being the limit of an increasing sequence of Riemann integrable functions.