



planetmath.org

Math for the people, by the people.

equivalent formulations for continuity

Canonical name	EquivalentFormulationsForContinuity
Date of creation	2013-03-22 15:18:23
Last modified on	2013-03-22 15:18:23
Owner	matte (1858)
Last modified by	matte (1858)
Numerical id	14
Author	matte (1858)
Entry type	Theorem
Classification	msc 26A15
Classification	msc 54C05
Related topic	Characterization

Suppose $f: X \rightarrow Y$ is a function between topological spaces X, Y . Then the following are equivalent:

1. f is continuous.
2. If B is open in Y , then $f^{-1}(B)$ is open in X .
3. If B is closed in Y , then $f^{-1}(B)$ is closed in X .
4. $f(\overline{A}) \subseteq \overline{f(A)}$ for all $A \subseteq X$.
5. If (x_i) is a net in X converging to x , then $(f(x_i))$ is a net in Y converging to $f(x)$. The concept of net can be replaced by the more familiar one of sequence if the spaces X and Y are first countable.
6. Whenever two nets S and T in X converge to the same point, then $f \circ S$ and $f \circ T$ converge to the same point in Y .
7. If \mathbb{F} is a filter on X that converges to x , then $f(\mathbb{F})$ is a filter on Y that converges to $f(x)$. Here, $f(\mathbb{F})$ is the filter generated by the filter base $\{f(F) \mid F \in \mathbb{F}\}$.
8. If B is any element of a <http://planetmath.org/Subbasis> \mathcal{S} for the topology of Y , then $f^{-1}(B)$ is open in X .
9. If B is any element of a basis \mathcal{B} for the topology of Y , then $f^{-1}(B)$ is open in X .
10. If $x \in X$, and N is any neighborhood of $f(x)$, then $f^{-1}(N)$ is a neighborhood of x .
11. f is continuous at every point in X .