

planetmath.org

Math for the people, by the people.

existence of nth root

Canonical name ExistenceOfNthRoot
Date of creation 2013-03-22 15:52:15
Last modified on Owner Wkbj79 (1863)
Last modified by Wkbj79 (1863)

Numerical id 21

Author Wkbj79 (1863)

Entry type Theorem
Classification msc 26C10
Classification msc 26A06
Classification msc 12D99

Related topic ExistenceOfNthRoot

Theorem. If $a \in \mathbb{R}$ with a > 0 and n is a positive integer, then there exists a unique positive real number u such that $u^n = a$.

Proof. The statement is clearly true for n = 1 (let u = a). Thus, it will be assumed that n > 1.

Define $p: \mathbb{R} \to \mathbb{R}$ by $p(x) = x^n - a$. Note that a positive real root of p(x) corresponds to a positive real number u such that $u^n = a$.

If a = 1, then $p(1) = 1^n - 1 = 0$, in which case the existence of u has been established.

Note that p(x) is a polynomial function and thus is continuous. If a < 1, then $p(1) = 1^n - a > 1 - 1 = 0$. If a > 1, then $p(a) = a^n - a = a(a^{n-1} - 1) > 0$. Note also that $p(0) = 0^n - a = -a < 0$. Thus, if $a \ne 1$, then the intermediate value theorem can be applied to yield the existence of u.

For uniqueness, note that the function p(x) is strictly increasing on the interval $(0, \infty)$. It follows that u as described in the statement of the theorem exists uniquely.