



planetmath.org

Math for the people, by the people.

nabla

Canonical name	Nabla
Date of creation	2013-03-22 14:00:20
Last modified on	2013-03-22 14:00:20
Owner	stevecheng (10074)
Last modified by	stevecheng (10074)
Numerical id	7
Author	stevecheng (10074)
Entry type	Definition
Classification	msc 26A06
Related topic	gradient
Related topic	NablaActingOnProducts
Related topic	Gradient
Related topic	AlternateCharacterizationOfCurl
Defines	∇

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a $C^1(\mathbb{R}^n)$ function, that is, a partially differentiable function in all its coordinates. The symbol ∇ , named *nabla*, represents the gradient operator, whose action on $f(x_1, x_2, \dots, x_n)$ is given by

$$\begin{aligned}\nabla f &= (f_{x_1}, f_{x_2}, \dots, f_{x_n}) \\ &= \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)\end{aligned}$$

Properties

1. If f, g are functions, then

$$\nabla(fg) = (\nabla f)g + f\nabla g.$$

2. For any scalars α and β and functions f and g ,

$$\nabla(\alpha f + \beta g) = \alpha \nabla f + \beta \nabla g.$$

The ∇ symbolism

Using the ∇ formalism, the divergence operator can be expressed as $\nabla \cdot$, the curl operator as $\nabla \times$, and the Laplacian operator as ∇^2 . To wit, for a given vector field

$$\mathbf{A} = A_x \mathbf{i} + A_y \mathbf{j} + A_z \mathbf{k},$$

and a given function f we have

$$\begin{aligned}\nabla \cdot \mathbf{A} &= \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \\ \nabla \times \mathbf{A} &= \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) \mathbf{i} + \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) \mathbf{j} + \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) \mathbf{k} \\ \nabla^2 f &= \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}.\end{aligned}$$