



planetmath.org

Math for the people, by the people.

Landau notation

Canonical name	LandauNotation
Date of creation	2013-03-22 11:42:56
Last modified on	2013-03-22 11:42:56
Owner	Mathprof (13753)
Last modified by	Mathprof (13753)
Numerical id	28
Author	Mathprof (13753)
Entry type	Definition
Classification	msc 26A12
Classification	msc 20H15
Classification	msc 20B30
Synonym	O notation
Synonym	omega notation
Synonym	theta notation
Synonym	big-O notation
Related topic	LowerBoundForSorting
Related topic	ConvergenceOfIntegrals
Defines	big-o
Defines	small-o
Defines	small-omega

Given two functions f and g from \mathbb{R}^+ to \mathbb{R}^+ , the notation

$$f = O(g)$$

means that the ratio $\frac{f(x)}{g(x)}$ stays bounded as $x \rightarrow \infty$. If moreover that ratio approaches zero, we write

$$f = o(g).$$

It is legitimate to write, say, $2x = O(x) = O(x^2)$, with the understanding that we are using the equality sign in an unsymmetric (and informal) way, in that we do not have, for example, $O(x^2) = O(x)$.

The notation

$$f = \Omega(g)$$

means that the ratio $\frac{f(x)}{g(x)}$ is bounded away from zero as $x \rightarrow \infty$, or equivalently $g = O(f)$.

If both $f = O(g)$ and $f = \Omega(g)$, we write $f = \Theta(g)$.

One more notational convention in this group is

$$f(x) \sim g(x),$$

meaning $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 1$.

In analysis, such notation is useful in describing error <http://planetmath.org/AsymptoticEst>. For example, the Riemann hypothesis is equivalent to the conjecture

$$\pi(x) = \text{li } x + O(\sqrt{x} \log x),$$

where $\text{li } x$ denotes the logarithmic integral.

Landau notation is also handy in applied mathematics, e.g. in describing the time complexity of an algorithm. It is common to say that an algorithm requires $O(x^3)$ steps, for example, without needing to specify exactly what is a step; for if $f = O(x^3)$, then $f = O(Ax^3)$ for any positive constant A .