



Math for the people, by the people.

generalisation of Gaussian integral

Canonical name	GeneralisationOfGaussianIntegral
Date of creation	2013-03-22 18:43:36
Last modified on	2013-03-22 18:43:36
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	6
Author	pahio (2872)
Entry type	Derivation
Classification	msc 26B15
Classification	msc 26A36
Related topic	SubstitutionNotation

The integral

$$\int_0^\infty e^{-x^2} \cos tx \, dx := w(t)$$

is a generalisation of the Gaussian integral $w(0) = \frac{\sqrt{\pi}}{2}$. For evaluating it we first form its derivative which may be done by <http://planetmath.org/DifferentiationUnderIntegral> under the integral sign:

$$w'(t) = \int_0^\infty e^{-x^2} (-x) \sin tx \, dx = \frac{1}{2} \int_0^\infty e^{-x^2} (-2x) \sin tx \, dx$$

Using integration by parts this yields

$$w'(t) = \frac{1}{2} \int_0^\infty e^{-x^2} \sin tx - \frac{t}{2} \int_0^\infty e^{-x^2} \cos tx \, dx = \frac{1}{2}(0-0) - \frac{t}{2} \int_0^\infty e^{-x^2} \cos tx \, dx = -\frac{t}{2}w(t).$$

Thus $w(t)$ satisfies the linear differential equation

$$\frac{dw}{dt} = -\frac{1}{2}tw,$$

where one can <http://planetmath.org/SeparationOfVariables> separate the variables and integrate:

$$\int \frac{dw}{w} = -\frac{1}{2} \int t \, dt.$$

So, $\ln w = -\frac{1}{4}t^2 + \ln C$, i.e. $w = w(t) = Ce^{-\frac{1}{4}t^2}$, and since there is the initial condition $w(0) = \frac{\sqrt{\pi}}{2}$, we obtain the result

$$w(t) = \frac{\sqrt{\pi}}{2}e^{-\frac{1}{4}t^2}.$$