

## general formulas for integration

Canonical name GeneralFormulasForIntegration

Date of creation 2013-03-22 17:39:31 Last modified on 2013-03-22 17:39:31

Owner pahio (2872) Last modified by pahio (2872)

Numerical id 16

Author pahio (2872)

Entry type Topic

Classification msc 26A36

Synonym integration formulas
Related topic TableOfDerivatives
Related topic IntegralTables

Related topic IntegrationByParts

Related topic ReductionFormulasForIntegrationOfPowers

$$1. \int f'(x) \, dx = f(x) + C$$

$$2. \int \lambda \, dx = \lambda x + C$$

3. 
$$\int \lambda f(x) \, dx = \lambda \int f(x) \, dx$$

4. 
$$\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$$

5. 
$$\int f(x)g'(x) dx = f(x)g(x) - \int g(x)f'(x) dx$$

6. 
$$\int g(f(x))f'(x) dx = G(f(x)) + C$$
 if  $G'(t) = g(t)$ 

7. 
$$\int [f(x)]^r f'(x) dx = \frac{1}{r+1} [f(x)]^{r+1} + C$$
 for  $r \neq -1$ 

8. 
$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + C$$

9. 
$$\int e^{f(x)} f'(x) dx = e^{f(x)} + C$$

10. 
$$\int \frac{f(x)}{(f(x)+a)(f(x)+b)} dx = \frac{a}{a-b} \int \frac{dx}{f(x)+a} - \frac{b}{a-b} \int \frac{dx}{f(x)+b}$$

11. 
$$\int \sin(\omega x + \varphi) dx = -\frac{\cos(\omega x + \varphi)}{\omega} + C$$

12. 
$$\int \cos(\omega x + \varphi) dx = \frac{\sin(\omega x + \varphi)}{\omega} + C$$

13. 
$$\int \sinh(\omega x + \varphi) dx = \frac{\cosh(\omega x + \varphi)}{\omega} + C$$

14. 
$$\int \cosh(\omega x + \varphi) dx = \frac{\sinh(\omega x + \varphi)}{\omega} + C$$

15. 
$$\int \sqrt{ax+b} \ dx = \frac{2}{3a}(ax+b)\sqrt{ax+b} + C$$

16. 
$$\int \sqrt{ax^2 + b} \, dx = \frac{x}{2} \sqrt{ax^2 + b} + \frac{b}{2\sqrt{a}} \ln(x\sqrt{a} + \sqrt{ax^2 + b}) + C$$

17. 
$$\int \sin^n x \cos^m x \, dx = -\frac{\sin^{n-1} x \cos^{m+1} x}{m+n} + \frac{n-1}{m+n} \int \sin^{n-2} x \cos^m x \, dx$$

18. 
$$\int \sin^n x \cos^m x \, dx = \frac{\sin^{n+1} x \cos^{m-1} x}{m+n} + \frac{m-1}{m+n} \int \sin^n x \cos^{m-2} x \, dx$$

Some series-formed antiderivatives: 
$$\int f(x) dx = C + f(0)x + \frac{f'(0)}{2!}x^2 + \frac{f''(0)}{3!}x^3 + \dots$$
$$\int f(x) dx = C + xf(x) - \frac{x^2}{2!}f'(x) + \frac{x^3}{3!}f''(x) - + \dots$$
$$\int UV dx = UV^{(-1)} - U'V^{(-2)} + U''V^{(-3)} - + \dots = \sum_{n=0}^{\infty} (-1)^n U^{(n)}V^{(-n-1)}$$

The derivatives with negative http://planetmath.org/HigherOrderDerivativesorder that V has been integrated repeatedly.