



planetmath.org

Math for the people, by the people.

chain rule

| | |
|------------------|-------------------------------------|
| Canonical name | ChainRule |
| Date of creation | 2013-03-22 12:26:43 |
| Last modified on | 2013-03-22 12:26:43 |
| Owner | matte (1858) |
| Last modified by | matte (1858) |
| Numerical id | 12 |
| Author | matte (1858) |
| Entry type | Theorem |
| Classification | msc 26A06 |
| Related topic | Derivative |
| Related topic | ChainRuleSeveralVariables |
| Related topic | ExampleOnSolvingAFunctionalEquation |
| Related topic | GudermannianFunction |

Let f, g be differentiable, real-valued functions such that g is defined on an open set $I \subseteq \mathbb{R}$, and f is defined on $g(I)$. Then the derivative of the composition $f \circ g$ is given by the *chain rule*, which asserts that

$$(f \circ g)'(x) = (f' \circ g)(x) g'(x), \quad x \in I.$$

The chain rule has a particularly suggestive appearance in terms of the Leibniz formalism. Suppose that z depends differentiably on y , and that y in turn depends differentiably on x . Then we have

$$\frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx}.$$

The apparent cancellation of the dy term is at best a formal mnemonic, and does not constitute a rigorous proof of this result. Rather, the Leibniz format is well suited to the interpretation of the chain rule in terms of related rates. To wit:

The instantaneous rate of change of z relative to x is equal to the rate of change of z relative to y times the rate of change of y relative to x .