



planetmath.org

Math for the people, by the people.

Hessian form

Canonical name	HessianForm
Date of creation	2013-03-22 15:00:03
Last modified on	2013-03-22 15:00:03
Owner	PrimeFan (13766)
Last modified by	PrimeFan (13766)
Numerical id	6
Author	PrimeFan (13766)
Entry type	Definition
Classification	msc 26B12
Related topic	HessianMatrix
Related topic	RelationsBetweenHessianMatrixAndLocalExtrema
Related topic	TestsForLocalExtremaForLagrangeMultiplierMethod

Given a smooth manifold M and $f: M \rightarrow \mathbb{R}$ being in $C^2(M)$, if x is a critical point of f , that is $df = 0$ at x , then we can define a symmetric 2-form

$$H(u_x, v_x) = u(v(f)) = v(u(f)),$$

where $H \in T^{*\otimes 2}$ and u and v are any vector fields that take the values u_x and v_x , respectively, at point x . Equality of the two defining expressions follows from the fact that x is a critical point of f , because then $[u, v](f) = df([u, v]) = 0$, where $[u, v]$ denotes the Lie bracket of the two vector fields. The form H is called the *Hessian form*.

In local coordinates, the Hessian form is given by

$$H = \frac{\partial^2 f}{\partial x^i \partial x^j} dx^i \otimes dx^j.$$

Its components are those of the Hessian matrix in the same coordinates. The advantage of the above formulation is coordinate independence. However, the price is that the Hessian form is only defined at critical points. It does not define a tensor field as one would naïvely expect.

Using the Hessian form, it is possible to analyze the critical points of f (determine whether they are local minima, maxima, or saddle points) in a coordinate independent way.