



planetmath.org

Math for the people, by the people.

## Euler's theorem on homogeneous functions

Canonical name	EulersTheoremOnHomogeneousFunctions
Date of creation	2013-03-22 15:18:58
Last modified on	2013-03-22 15:18:58
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	10
Author	CWoo (3771)
Entry type	Theorem
Classification	msc 26B12
Classification	msc 26A06
Classification	msc 15-00
Defines	Euler operator

**Theorem 1** (Euler). *Let  $f(x_1, \dots, x_k)$  be a smooth homogeneous function of degree  $n$ . That is,*

$$f(tx_1, \dots, tx_k) = t^n f(x_1, \dots, x_k). \quad (*)$$

*Then the following identity holds*

$$x_1 \frac{\partial f}{\partial x_1} + \dots + x_k \frac{\partial f}{\partial x_k} = n f.$$

*Proof.* By homogeneity, the relation (??) holds for all  $t$ . Taking the  $t$ -derivative of both sides, we establish that the following identity holds for all  $t$ :

$$x_1 \frac{\partial f}{\partial x_1}(tx_1, \dots, tx_k) + \dots + x_k \frac{\partial f}{\partial x_k}(tx_1, \dots, tx_k) = nt^{n-1} f(x_1, \dots, x_k).$$

To obtain the result of the theorem, it suffices to set  $t = 1$  in the previous formula.  $\square$

Sometimes the differential operator  $x_1 \frac{\partial}{\partial x_1} + \dots + x_k \frac{\partial}{\partial x_k}$  is called the *Euler operator*. An equivalent way to state the theorem is to say that homogeneous functions are eigenfunctions of the Euler operator, with the degree of homogeneity as the eigenvalue.