

# planetmath.org

Math for the people, by the people.

#### table of derivatives

Canonical name TableOfDerivatives
Date of creation 2013-03-22 17:34:48
Last modified on 2013-03-22 17:34:48

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 31

Author CWoo (3771)

Entry type Feature
Classification msc 26A24
Related topic Integral Tables
Related topic Derivative2

Related topic GeneralFormulasForIntegration

#### Basic rules

f(x)	$\frac{df(x)}{dx} = f'(x)$
f(x) + g(x)	f'(x) + g'(x)
f(x)g(x)	f'(x)g(x) + f(x)g'(x)
$f(x) \over g(x), g \neq 0$	$\frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$
f(g(x))	f'(g(x))g'(x)
$f^{-1}(x)$	$\frac{1}{f'(f^{-1}(x))}$

 $\verb|http://planetmath.org/Polynomial* Polynomials and powers|$ 

f(x)	f'(x)	applicable domain
$c \in \mathbb{R}$	0	$x \in \mathbb{R}$
$x^r$	$rx^{r-1}$	$x \in \mathbb{R}$
$\sqrt{x}$	$\frac{1}{2\sqrt{x}}$	x > 0
	$\frac{x}{ x } = \frac{ x }{x}$	$x \neq 0$

### Exponential and logarithmic functions

f(x)	f'(x)	applicable domain
$\exp(x) = e^x$	$\exp(x) = e^x$	$x \in \mathbb{R}$
$a^x$	$a^x \ln a$	$x \in \mathbb{R}$
$\ln x$	$\frac{1}{x}$	x > 0
$x^x$	$x^x(1+\ln x)$	x > 0

## $\verb|http://planetmath.org/Trigonometry Trigonometric functions|\\$

f(x)	f'(x)	applicable domain
$\sin x$	$\cos x$	$x \in \mathbb{R}$
$\cos x$	$-\sin x$	$x \in \mathbb{R}$
$\tan x$	$\sec^2 x$	$x \neq n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$
$\cot x$	$-\csc^2 x$	$x \neq n\pi, n \in \mathbb{Z}$
$\sec x$	$\sec x \tan x$	$x \neq n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$
$\csc x$	$-\csc x \cot x$	$x \neq n\pi, n \in \mathbb{Z}$
$\arcsin x$	$\frac{1}{\sqrt{1-x^2}}$	x  < 1
$\arccos x$	$-\frac{1}{\sqrt{1-x^2}}$	x  < 1
$\arctan x$	$\frac{1}{1+x^2}$	$x \in \mathbb{R}$

 $\verb|http://planetmath.org/HyperbolicFunctions| Hyperbolic functions|$ 

f(x)	f'(x)	applicable domain
$\sinh x$	$\cosh x$	$x \in \mathbb{R}$
$\cosh x$	$\sinh x$	$x \in \mathbb{R}$
$\tanh x$	$\operatorname{sech}^2 x$	$x \in \mathbb{R}$
$\coth x$	$-\operatorname{csch}^2 x$	$x \neq 0$
$\operatorname{sech} x$	$-\operatorname{sech} x \tanh x$	$x \in \mathbb{R}$
$\operatorname{csch} x$	$-\operatorname{csch} x \operatorname{coth} x$	$x \neq 0$
arsinh x	$\frac{1}{\sqrt{x^2+1}}$	$x \neq 0$
$\operatorname{arcosh} x$	$\frac{1}{\sqrt{x^2-1}}$	x  > 1
$\operatorname{artanh} x$	$\frac{1}{1-x^2}$	-1 < x < 1
$\operatorname{arcoth} x$	$\frac{1}{1-x^2}$	x  > 1

#### Other functions

(see error function, logarithmic integral, sine integral, Gudermannian, Hermite polynomials)  $\,$ 

f(x)	f'(x)	applicable domain
$\operatorname{Erf} x$	$\frac{2}{\sqrt{\pi}}e^{-x^2}$	$x \in \mathbb{R}$
Li x	$\frac{1}{\ln x}$	x > 1
$\operatorname{Si} x$	$\operatorname{sinc} x$	$x \in \mathbb{R}$
$\operatorname{gd} x$	$\frac{1}{\cosh x}$	$x \in \mathbb{R}$
$gd^{-1}x$	$\frac{1}{\cos x}$	$ x  < \frac{\pi}{2}$
$H_n(x)$	$2nH_{n-1}(x)$	$x \in \mathbb{R}$

Instructions on how to add a function and its derivative. Open the entry in edit mode. Using the appropriate table for your function (or make a new table if applicable), make a copy of the two lines of comment (starting with %) in the code (within the tabular environment) and paste it immediately before the comment. For functions outside of the "Basic rules" section, include the appropriate domain. Uncomment the lines (take out the % symbols) after completing. Preview before saving the entry.