



planetmath.org

Math for the people, by the people.

average value of function

Canonical name	AverageValueOfFunction
Date of creation	2013-03-22 19:01:54
Last modified on	2013-03-22 19:01:54
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	12
Author	pahio (2872)
Entry type	Definition
Classification	msc 26D15
Classification	msc 11-00
Related topic	ArithmeticMean
Related topic	Mean3
Related topic	Countable
Related topic	GaussMeanValueTheorem
Related topic	Expectation
Related topic	MeanSquareDeviation
Defines	average value

The set of the values of a real function  $f$  defined on an interval  $[a, b]$  is usually uncountable, and therefore for being able to speak of an *average value* of  $f$  in the sense of the average value

$$A.V. = \frac{a_1 + a_2 + \dots + a_n}{n} = \frac{\sum_{j=1}^n a_j}{\sum_{j=1}^n 1} \quad (1)$$

of a finite list  $a_1, a_2, \dots, a_n$  of numbers, one has to replace the sums with integrals. Thus one could define

$$A.V.(f) := \frac{\int_a^b f(x) dx}{\int_a^b 1 dx},$$

i.e.

$$A.V.(f) := \frac{1}{b-a} \int_a^b f(x) dx. \quad (2)$$

For example, the average value of  $x^2$  on the interval  $[0, 1]$  is  $\frac{1}{3}$  and the average value of  $\sin x$  on the interval  $[0, \pi]$  is  $\frac{2}{\pi}$ .

The definition (2) may be extended to complex functions  $f$  on an arc  $\gamma$  of a rectifiable curve via the contour integral

$$A.V.(f) := \frac{1}{l(\gamma)} \int_{\gamma} f(z) dz \quad (3)$$

where  $l(\gamma)$  is the <http://planetmath.org/ArcLength>length of the arc. If especially  $\gamma$  is a closed curve in a simply connected domain where  $f$  is analytic, then the average value of  $f$  on  $\gamma$  is always 0, as the Cauchy integral theorem implies.