



planetmath.org

Math for the people, by the people.

Jacobian matrix

Canonical name	JacobianMatrix
Date of creation	2013-03-22 11:58:33
Last modified on	2013-03-22 11:58:33
Owner	PhysBrain (974)
Last modified by	PhysBrain (974)
Numerical id	18
Author	PhysBrain (974)
Entry type	Definition
Classification	msc 26B10
Synonym	derivative of a multivariable function
Synonym	derivative of a vector-valued function
Synonym	Jacobi matrix
Related topic	PartialDerivative
Related topic	Derivative2
Related topic	DerivativeNotation
Related topic	Gradient
Related topic	ChainRuleSeveralVariables
Related topic	DirectionalDerivative
Related topic	JordanBanachAndJordanLieAlgebras
Defines	Jacobian

The *Jacobian matrix* $[\mathbf{J}f(\mathbf{a})]$ of a function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ at the point \mathbf{a} with respect to some choice of bases for \mathbb{R}^n and \mathbb{R}^m is the matrix of the linear map from \mathbb{R}^n into \mathbb{R}^m that generalizes the definition of the derivative of a function on \mathbb{R} . It can be defined as the matrix of the linear map D , such that

$$\lim_{\mathbf{h} \rightarrow \mathbf{0}} \frac{\|f(\mathbf{a} + \mathbf{h}) - f(\mathbf{a}) - D(\mathbf{h})\|}{\|\mathbf{h}\|} = 0$$

The linear map that satisfies the above limit is called the derivative of f at \mathbf{a} . It is easy to show that the Jacobian matrix of a given differentiable function at \mathbf{a} with respect to chosen bases is just the matrix of partial derivatives of the component functions of f at \mathbf{a} :

$$[\mathbf{J}f(\mathbf{x})] = \begin{bmatrix} D_1 f_1(\mathbf{x}) & \dots & D_n f_1(\mathbf{x}) \\ \vdots & \ddots & \vdots \\ D_1 f_m(\mathbf{x}) & \dots & D_n f_m(\mathbf{x}) \end{bmatrix}$$

A more concise way of writing it is

$$[\mathbf{J}f(\mathbf{x})] = [\overrightarrow{D_1 f}, \dots, \overrightarrow{D_n f}] = \begin{bmatrix} \nabla f_1 \\ \vdots \\ \nabla f_m \end{bmatrix}$$

where $\overrightarrow{D_n f}$ is the partial derivative with respect to the n 'th variable and ∇f_m is the gradient of the m 'th component of \mathbf{f} .

Note that the Jacobian matrix represents the matrix of the derivative D of f at \mathbf{x} iff f is differentiable at \mathbf{x} . Also, if f is differentiable at \mathbf{x} , then the directional derivative in the direction \vec{v} is $D(\vec{v}) = [\mathbf{J}f(\mathbf{x})]\vec{v}$.

Given local coordinates for some real submanifold M of \mathbb{R}^n , it is easy to show that the effect of a change of coordinates on volume forms is a local scaling of the volume form by the determinant of the Jacobian matrix of the derivative of the backwards change of coordinates, which is called the inverse Jacobian. The determinant of the inverse Jacobian is thus commonly seen in integration over a change of coordinates.

$$\int_{\Omega_x} f(x) d\Omega_x = \int_{\Omega_\xi} f(\xi) |J^{-1}| d\Omega_\xi$$

where $x \in \Omega_x \subset M$, $\xi \in \Omega_\xi \subset M$, and $\mathbf{J} = \nabla \xi(x)$ is the derivative of the function $\xi(x)$ which maps points in Ω_x to points in Ω_ξ .