

## alternative proof of derivative of $x^n$

Canonical name AlternativeProofOfDerivativeOfXn

Date of creation 2013-03-22 15:59:31 Last modified on 2013-03-22 15:59:31 Owner Wkbj79 (1863) Last modified by Wkbj79 (1863)

Numerical id 12

Author Wkbj79 (1863)

Entry type Proof

Classification msc 26B05 Classification msc 26A24 Related topic DerivativeOfXn

Related topic DerivativesByPureAlgebra

The typical derivative formula

$$\frac{df}{dx} = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

combined with the binomial theorem yield an alternative way to prove that

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

for any positive integer n.

$$\frac{d}{dx}(x^n) = \lim_{h \to 0} \frac{(x+h)^n - x^n}{h}$$

$$= \lim_{h \to 0} \frac{\left(\sum_{j=0}^n \binom{n}{j} x^j h^{n-j}\right) - x^n}{h}$$

Proof. 
$$x^{n} + nx^{n-1}h + h^{2}\left(\sum_{j=0}^{n-2} \binom{n}{j}x^{j}h^{n-2-j}\right) - x^{n}$$

$$= \lim_{h \to 0} \frac{1}{h}$$

$$=\lim_{h\to 0}\frac{nx^{n-1}h+h^2\displaystyle\sum_{j=0}^{n-2}\binom{n}{j}x^jh^{n-2-j}}{h}$$

$$= \lim_{h \to 0} \left( nx^{n-1} + h \sum_{j=0}^{n-2} \binom{n}{j} x^j h^{n-2-j} \right)$$

$$= nx^{n-1}$$