



planetmath.org

Math for the people, by the people.

fundamental theorems of calculus for  
Lebesgue integration

Canonical name	FundamentalTheoremsOfCalculusForLebesgueIntegration
Date of creation	2013-03-22 12:27:54
Last modified on	2013-03-22 12:27:54
Owner	mathcam (2727)
Last modified by	mathcam (2727)
Numerical id	17
Author	mathcam (2727)
Entry type	Theorem
Classification	msc 26-00
Synonym	first fundamental theorem of calculus
Synonym	second fundamental theorem of calculus
Synonym	fundamental theorem of calculus
Related topic	FundamentalTheoremOfCalculusClassicalVersion
Related topic	FundamentalTheoremOfCalculusForRiemannIntegration
Related topic	ChangeOfVariableInDefiniteIntegral

Loosely, the *Fundamental Theorems of Calculus* serve to demonstrate that integration and differentiation are inverse processes. Suppose that  $F(x)$  is an absolutely continuous function on an interval  $[a, b] \subset \mathbb{R}$ . The two following forms of the theorem are equivalent.

**First form of the Fundamental Theorem:**

There exists a function  $f(t)$  Lebesgue-integrable on  $[a, b]$  such that for any  $x \in [a, b]$ , we have  $F(x) - F(a) = \int_a^x f(t)dt$ .

**Second form of the Fundamental Theorem:**

$F(x)$  is differentiable almost everywhere on  $[a, b]$  and its derivative, denoted  $F'(x)$ , is Lebesgue-integrable on that interval. In addition, we have the relation  $F(x) - F(a) = \int_a^x F'(t)dt$  for any  $x \in [a, b]$ .