



planetmath.org

Math for the people, by the people.

facts about Riemann–Stieltjes integral

Canonical name	FactsAboutRiemannStieltjesIntegral
Date of creation	2013-03-22 18:55:03
Last modified on	2013-03-22 18:55:03
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	5
Author	pahio (2872)
Entry type	Topic
Classification	msc 26A42
Synonym	properties of Riemann–Stieltjes integral
Related topic	PropertiesOfRiemannStieltjesIntegral

- If the integrator g of the <http://planetmath.org/node/3187>Riemann–Stieltjes integral $\int_a^b f(x) dg(x)$ is the identity function, then the integral reduces to the Riemann integral $\int_a^b f(x) dx$.

- If the integrand of the Riemann–Stieltjes integral is a constant function, one has

$$\int_a^b c dg(x) = c \cdot (g(b) - g(a)).$$

- If the integrand f is continuous and the integrator g monotonically nondecreasing on the interval $[a, b]$, then there exists a number ξ on the interval such that

$$\int_a^b f(x) dg(x) = f(\xi)(g(b) - g(a)).$$

Cf. the integral mean value theorem.

- If f is continuous, g monotonically nondecreasing and differentiable on the interval $[a, b]$, then

$$\frac{d}{dx} \int_a^x f(t) dg(t) = f(x)g'(x) \quad \text{for } a < x < b.$$