



planetmath.org

Math for the people, by the people.

gradient theorem

Canonical name	GradientTheorem
Date of creation	2013-03-22 19:11:16
Last modified on	2013-03-22 19:11:16
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	8
Author	pahio (2872)
Entry type	Theorem
Classification	msc 26B12
Synonym	fundamental theorem of line integrals
Related topic	LaminarField
Related topic	Gradient

If $u = u(x, y, z)$ is continuously differentiable function in a simply connected <http://planetmath.org/Domain2domain> D of \mathbb{R}^3 and $P = (x_0, y_0, z_0)$ and $Q = (x_1, y_1, z_1)$ lie in this domain, then

$$\int_P^Q \nabla u \cdot \vec{ds} = u(x_1, y_1, z_1) - u(x_0, y_0, z_0) \quad (1)$$

where the line integral of the left hand side is taken along an arbitrary path in D .

The equation (1) is illustrated by the fact that

$$\nabla u \cdot \vec{ds} = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy + \frac{\partial u}{\partial z} dz$$

is the total differential of u , and thus

$$\int_P^Q \nabla u \cdot \vec{ds} = \int_P^Q du.$$