



planetmath.org

Math for the people, by the people.

Hardy-Littlewood maximal operator

Canonical name	HardyLittlewoodMaximalOperator
Date of creation	2013-03-22 13:27:30
Last modified on	2013-03-22 13:27:30
Owner	azdbacks4234 (14155)
Last modified by	azdbacks4234 (14155)
Numerical id	8
Author	azdbacks4234 (14155)
Entry type	Definition
Classification	msc 28A25
Classification	msc 28A15
Related topic	HardyLittlewoodMaximalTheorem
Defines	Hardy-Littlewood maximal function

The *Hardy-Littlewood maximal operator* in \mathbb{R}^n is an operator defined on $L^1_{\text{loc}}(\mathbb{R}^n)$ (the space of locally integrable functions in \mathbb{R}^n with the Lebesgue measure) which maps each locally integrable function f to another function Mf , defined for each $x \in \mathbb{R}^n$ by

$$Mf(x) = \sup_Q \frac{1}{m(Q)} \int_Q |f(y)| dy,$$

where the supremum is taken over all cubes Q containing x . This function is lower semicontinuous (and hence measurable), and it is called the *Hardy-Littlewood maximal function* of f .

The operator M is sublinear, which means that

$$M(af + bg) \leq |a|Mf + |b|Mg$$

for each pair of locally integrable functions f, g and scalars a, b .