



Math for the people, by the people.

signed measure

Canonical name	SignedMeasure
Date of creation	2013-03-22 13:26:55
Last modified on	2013-03-22 13:26:55
Owner	Koro (127)
Last modified by	Koro (127)
Numerical id	8
Author	Koro (127)
Entry type	Definition
Classification	msc 28A12

A *signed measure* on a measurable space  $(\Omega, \mathcal{S})$  is a function  $\mu : \mathcal{S} \rightarrow \mathbb{R} \cup \{+\infty\}$  which is [`http://planetmath.org/Additive`](http://planetmath.org/Additive) $\sigma$ -additive and such that  $\mu(\emptyset) = 0$ .

**Remarks.**

1. The usual (positive) measure is a particular case of signed measure, in which  $|\mu| = \mu$  (see Jordan decomposition.)
2. Notice that the value  $-\infty$  is not allowed. For some authors, a signed measure can only take finite values (so that  $+\infty$  is not allowed either). This is sometimes useful because it turns the space of all signed measures into a normed vector space, with the natural operations, and the norm given by  $\|\mu\| = |\mu|(\Omega)$ .
3. An important example of signed measures arises from the usual measures in the following way: Let  $(\Omega, \mathcal{S}, \mu)$  be a measure space, and let  $f$  be a (real valued) measurable function such that

$$\int_{\{x \in \Omega : f(x) < 0\}} |f| d\mu < \infty.$$

Then a signed measure is defined by

$$A \mapsto \int_A f d\mu.$$