



planetmath.org

Math for the people, by the people.

Jordan decomposition

| | |
|------------------|---------------------|
| Canonical name | JordanDecomposition |
| Date of creation | 2013-03-22 13:27:02 |
| Last modified on | 2013-03-22 13:27:02 |
| Owner | Koro (127) |
| Last modified by | Koro (127) |
| Numerical id | 9 |
| Author | Koro (127) |
| Entry type | Definition |
| Classification | msc 28A12 |
| Defines | positive variation |
| Defines | negative variation |
| Defines | total variation |

Let $(\Omega, \mathcal{S}, \mu)$ be a signed measure space, and let (A, B) be a Hahn decomposition for μ . We define μ^+ and μ^- by

$$\mu^+(E) = \mu(A \cap E) \quad \text{and} \quad \mu^-(E) = -\mu(B \cap E).$$

This definition is easily shown to be independent of the chosen Hahn decomposition.

It is clear that μ^+ is a positive measure, and it is called the *positive variation* of μ . On the other hand, μ^- is a positive finite measure, called the *negative variation* of μ . The measure $|\mu| = \mu^+ + \mu^-$ is called the *total variation* of μ .

Notice that $\mu = \mu^+ - \mu^-$. This decomposition of μ into its positive and negative parts is called the *Jordan decomposition* of μ .