



Math for the people, by the people.

## Simpson's rule

Canonical name	SimpsonsRule
Date of creation	2013-03-22 13:40:12
Last modified on	2013-03-22 13:40:12
Owner	drini (3)
Last modified by	drini (3)
Numerical id	9
Author	drini (3)
Entry type	Theorem
Classification	msc 28-00
Classification	msc 26A06
Classification	msc 41A55
Classification	msc 65D32
Related topic	LagrangeInterpolationFormula
Related topic	NewtonAndCotesFormulas
Related topic	Prismatoid

*Simpson's rule* is a method of (approximate) numerical definite integration (or quadrature). Simpson's rule is based on a parabolic model of the function to be integrated (in contrast to the trapezoidal model of the trapezoidal rule). Thus, a minimum of three points and three function values are required. Here we take three equidistant points:  $x_0, x_2$  the interval endpoints,  $x_1 = (x_0 + x_2)/2$  the midpoint, and let  $h = |b - a|/2$  the distance between each. The definite integral is then approximated by:

$$\int_{x_0}^{x_2} f(x)dx \approx I = \frac{h}{3}(f(x_0) + 4f(x_1) + f(x_2))$$

We can extend this to greater precision by breaking our target domain into  $n$  equal-length fragments. The quadrature is then the weighted sum of the above formula for every pair of adjacent regions, which works out for even  $n$  to

$$I = \frac{h}{3}(f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \cdots + 4f(x_{n-3}) + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n))$$