



planetmath.org

Math for the people, by the people.

**continuous almost everywhere versus equal to
a continuous function almost everywhere**

Canonical name	ContinuousAlmostEverywhereVersusEqualToAContinuousFunctionAlmostEve
Date of creation	2013-03-22 15:58:47
Last modified on	2013-03-22 15:58:47
Owner	Wkbj79 (1863)
Last modified by	Wkbj79 (1863)
Numerical id	10
Author	Wkbj79 (1863)
Entry type	Example
Classification	msc 28A12
Classification	msc 60A10

The concept of almost everywhere can be somewhat tricky to people who are not familiar with it. Let m denote Lebesgue measure. Consider the following two statements about a function $f: \mathbb{R} \rightarrow \mathbb{R}$:

- f is <http://planetmath.org/Continuous> almost everywhere with respect to m
- f is equal to a continuous function almost everywhere with respect to m

Although these two statements seem alike, they have quite different meanings. In fact, neither one of these statements implies the other.

Consider the function $\chi_{[0,\infty)}(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0. \end{cases}$

This function is not continuous at 0, but it is continuous at all other $x \in \mathbb{R}$. Note that $m(\{0\}) = 0$. Thus, $\chi_{[0,\infty)}$ is continuous almost everywhere.

Suppose $\chi_{[0,\infty)}$ is equal to a continuous function almost everywhere. Let $A \subset \mathbb{R}$ be <http://planetmath.org/LebesgueMeasure> Lebesgue measurable with $m(A) = 0$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ such that $\chi_{[0,\infty)}(x) = g(x)$ for all $x \in \mathbb{R} \setminus A$. Since $\chi_{[0,\infty)}(x) = 0$ for all $x < 0$ and $m(A \cap (-\infty, 0)) = 0$, there exists $a < 0$ such that $g(a) = 0$. Similarly, there exists $b \geq 0$ such that $g(b) = 1$. Since g is continuous, by the intermediate value theorem, there exists $c \in (a, b)$ with $g(c) = \frac{1}{2}$. Let $U = (0, 1)$. Since g is continuous, $g^{-1}(U)$ is open. Recall that $c \in g^{-1}(U)$. Thus, $g^{-1}(U) \neq \emptyset$. Since $g^{-1}(U)$ is a nonempty open set, $m(g^{-1}(U)) > 0$. On the other hand, $g^{-1}(U) \subseteq A$, yielding that $0 < m(g^{-1}(U)) \leq m(A) = 0$, a contradiction.

Now consider the function $\chi_{\mathbb{Q}}(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q}. \end{cases}$

Note that $m(\mathbb{Q}) = 0$. Thus, $\chi_{\mathbb{Q}} = 0$ almost everywhere. Since 0 is continuous, $\chi_{\mathbb{Q}}$ is equal to a continuous function almost everywhere. On the other hand, $\chi_{\mathbb{Q}}$ is not continuous almost everywhere. Actually, $\chi_{\mathbb{Q}}$ is not continuous at any $x \in \mathbb{R}$. Recall that \mathbb{Q} and $\mathbb{R} \setminus \mathbb{Q}$ are both <http://planetmath.org/Dense> dense in \mathbb{R} . Therefore, for every $x \in \mathbb{R}$ and for every $\delta > 0$, there exist $x_1 \in (x - \delta, x + \delta) \cap \mathbb{Q}$ and $x_2 \in (x - \delta, x + \delta) \cap (\mathbb{R} \setminus \mathbb{Q})$. Since $\chi_{\mathbb{Q}}(x_1) = 1$ and $\chi_{\mathbb{Q}}(x_2) = 0$, it follows that $\chi_{\mathbb{Q}}$ is not continuous at x . (Choose any $\varepsilon \in (0, 1)$.)