



planetmath.org

Math for the people, by the people.

extending a capacity to a Cartesian product

Canonical name	ExtendingACapacityToACartesianProduct
Date of creation	2013-03-22 18:47:38
Last modified on	2013-03-22 18:47:38
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	6
Author	gel (22282)
Entry type	Theorem
Classification	msc 28A12
Classification	msc 28A05

A capacity on a set X can be extended to a set function on a Cartesian product $X \times K$ simply by projecting any subset onto X , and then applying the original capacity.

Theorem. *Suppose that (X, \mathcal{F}) is a paved space such that \mathcal{F} is closed under finite unions and finite intersections, and that (K, \mathcal{K}) is a compact paved space. Define \mathcal{G} to be the closure under finite unions and finite intersections of the paving $\mathcal{F} \times \mathcal{K}$ on $X \times K$.*

If I is an \mathcal{F} -capacity and $\pi_X: X \times K \rightarrow X$ is the projection map, we can form the composition

$$\begin{aligned} I \circ \pi_X: \mathcal{P}(X \times K) &\rightarrow \mathbb{R}, \\ I \circ \pi_X(S) &= I(\pi_X(S)). \end{aligned}$$

Then $\pi_X(S) \in \mathcal{F}_\delta$ for any $S \in \mathcal{G}_\delta$, and $I \circ \pi_X$ is a \mathcal{G} -capacity.

This result justifies looking at capacities when considering projections from the Cartesian product $X \times K$ onto X . We see that the property of being a capacity is preserved under composing with such projections. However, additivity of set functions is not preserved, so the corresponding result would not be true if “capacity” was replaced by “measure” or “outer measure”.

Recall that if $S \subseteq X \times K$ is $(\mathcal{G}, I \circ \pi_X)$ -capacitable then, for any $\epsilon > 0$, there is an $A \in \mathcal{G}_\delta$ such that $A \subseteq S$ and $I \circ \pi_X(A) > I \circ \pi_X(S) - \epsilon$. However, $\pi_X(A) \subseteq \pi_X(S)$ and, by the above theorem, $\pi_X(A) \in \mathcal{F}_\delta$. This has the following consequence.

Lemma. *Let $S \subseteq X \times K$ be $(\mathcal{G}, I \circ \pi_X)$ -capacitable. Then, $\pi_X(S)$ is (\mathcal{F}, I) -capacitable.*