



planetmath.org

Math for the people, by the people.

proof of equivalence of formulas for exp

Canonical name	ProofOfEquivalenceOfFormulasForExp
Date of creation	2013-03-22 15:22:52
Last modified on	2013-03-22 15:22:52
Owner	stevecheng (10074)
Last modified by	stevecheng (10074)
Numerical id	11
Author	stevecheng (10074)
Entry type	Proof
Classification	msc 30A99
Related topic	ComplexExponentialFunction
Related topic	ExponentialFunction
Related topic	MatrixExponential

We present an elementary proof that:

$$\sum_{k=0}^{\infty} \frac{z^k}{k!} = \lim_{n \rightarrow \infty} \left(1 + \frac{z}{n}\right)^n.$$

There are of course other proofs, but this one has the advantage that it carries verbatim for the matrix exponential and the operator exponential.

At the outset, we observe that  $\sum_{k=0}^{\infty} z^k/k!$  converges by the ratio test. For definiteness, the notation  $e^z$  below will refer to exactly this series.

*Proof.* We expand the right-hand in the straightforward manner:

$$\begin{aligned} \left(1 + \frac{z}{n}\right)^n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{z}{n}\right)^k \\ &= \sum_{k=0}^n \frac{n \cdot (n-1) \cdots (n-k+1)}{n^k} \frac{z^k}{k!} = \sum_{k=0}^n \pi(k, n) \frac{z^k}{k!}, \end{aligned}$$

where  $\pi(k, n)$  denotes the coefficient

$$1 \cdot \left(1 - \frac{1}{n}\right) \cdot \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{k-1}{n}\right).$$

Let  $|z| \leq M$ . Given  $\epsilon > 0$ , there is a  $N \in \mathbb{N}$  such that whenever  $n \geq N$ , then  $\sum_{k=n+1}^{\infty} M^k/k! < \epsilon/2$ , since the sum is the tail of the convergent series  $e^M$ .

Since  $\lim_{n \rightarrow \infty} \pi(k, n) = 1$  for  $k$ , there is also a  $N' \in \mathbb{N}$ , with  $N' \geq N$ , so that whenever  $n \geq N'$  and  $0 \leq k \leq N$ , then  $|\pi(k, n) - 1| < \epsilon/(2e^M)$ . (Note that  $k$  is chosen only from a *finite* set.)

Now, when  $n \geq N'$ , we have

$$\begin{aligned} \left| \sum_{k=0}^n \pi(k, n) \frac{z^k}{k!} - \sum_{k=0}^{\infty} \frac{z^k}{k!} \right| &= \left| \sum_{k=0}^n (\pi(k, n) - 1) \frac{z^k}{k!} - \sum_{k=n+1}^{\infty} \frac{z^k}{k!} \right| \\ &\leq \sum_{k=0}^n |\pi(k, n) - 1| \frac{M^k}{k!} + \sum_{k=n+1}^{\infty} \frac{M^k}{k!} \\ &= \sum_{k=0}^N |\pi(k, n) - 1| \frac{M^k}{k!} + \sum_{k=N+1}^n |\pi(k, n) - 1| \frac{M^k}{k!} + \sum_{k=n+1}^{\infty} \frac{M^k}{k!} \\ &< \frac{\epsilon}{2e^M} \sum_{k=0}^N \frac{M^k}{k!} + \sum_{k=N+1}^n \frac{M^k}{k!} + \sum_{k=n+1}^{\infty} \frac{M^k}{k!} \end{aligned}$$

(In the middle sum, we use the bound  $|\pi(k, n) - 1| = 1 - \pi(k, n) \leq 1$  for all  $k$  and  $n$ .)

$$< \frac{\epsilon}{2e^M} \cdot e^M + \frac{\epsilon}{2} = \epsilon. \quad \square$$

In fact, we have proved uniform convergence of  $\lim_{n \rightarrow \infty} \left(1 + \frac{z}{n}\right)^n$  over  $|z| \leq M$ . Exploiting this fact we can also show:

$$\left(1 + \frac{z}{n} + o\left(\frac{1}{n}\right)\right)^n = \left(1 + \frac{z + o(1)}{n}\right)^n \rightarrow \sum_{k=0}^{\infty} \frac{z^k}{k!} \quad (\text{pointwise, as } n \rightarrow \infty)$$

*Proof.*  $|z| < M$ . Given  $\epsilon > 0$ , for large enough  $n$ , we have

$$\left|\left(1 + \frac{w}{n}\right)^n - e^w\right| < \epsilon/2 \quad \text{uniformly for all } |w| \leq M.$$

Since  $o(1) \rightarrow 0$ , for large enough  $n$  we can set  $w = z + o(1)$  above. Since the exponential is continuous<sup>1</sup>, for large enough  $n$  we also have  $|e^{z+o(1)} - e^z| < \epsilon/2$ . Thus

$$\left|\left(1 + \frac{z + o(1)}{n}\right)^n - e^z\right| \leq \left|\left(1 + \frac{z + o(1)}{n}\right)^n - e^{z+o(1)}\right| + |e^{z+o(1)} - e^z| < \epsilon$$

---

<sup>1</sup>follows from uniform convergence on bounded subsets of either expression for  $e^z$