



planetmath.org

Math for the people, by the people.

holomorphic function associated with continuous function

Canonical name	HolomorphicFunctionAssociatedWithContinuousFunction
Date of creation	2013-03-22 19:14:29
Last modified on	2013-03-22 19:14:29
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	11
Author	pahio (2872)
Entry type	Theorem
Classification	msc 30E20
Classification	msc 30D20
Related topic	DifferentiationUnderIntegralSign
Related topic	CauchyIntegralFormula

Theorem. If $f(z)$ is continuous on a (finite) contour γ of the complex plane, then the contour integral

$$g(z) =: \int_{\gamma} \frac{f(t)}{t-z} dt, \quad (1)$$

defines a function $z \mapsto g(z)$ which is holomorphic in any domain D not containing points of γ . Moreover, the derivative has the expression

$$g'(z) = \int_{\gamma} \frac{f(t)}{(t-z)^2} dt. \quad (2)$$

Proof. The right hand side of (2) is defined since its integrand is continuous. One has to show that it equals

$$\lim_{\Delta z \rightarrow 0} \frac{g(z+\Delta z) - g(z)}{\Delta z}.$$

Let $z_1 =: z + \Delta z \notin \gamma$, $\Delta z \neq 0$. We may write first

$$\frac{g(z_1) - g(z)}{z_1 - z} = \frac{1}{\Delta z} \int_{\gamma} f(t) \left[\frac{1}{t-z_1} - \frac{1}{t-z} \right] dt = \int_{\gamma} \frac{f(t)}{(t-z_1)(t-z)} dt,$$

whence

$$E =: \frac{g(z_1) - g(z)}{z_1 - z} - \int_{\gamma} \frac{f(t)}{(t-z)^2} dt = \Delta z \cdot \int_{\gamma} \frac{f(t)}{(t-z_1)(t-z)^2} dt.$$

Because f is continuous in the compact set γ , there is a positive constant M such that

$$|f(t)| < M \quad \forall t \in \gamma.$$

As well, we have a positive constant d such that

$$|t-z| \geq d \quad \forall t \in \gamma.$$

When we choose $|\Delta z| < \frac{d}{2}$, it follows that

$$|t-z_1| = |(t-z) - \Delta z| \geq |t-z| - |\Delta z| > d - \frac{d}{2} = \frac{d}{2}.$$

Consequently,

$$\left| \frac{f(t)}{(t-z_1)(t-z)^2} \right| = \frac{|f(t)|}{|t-z_1||t-z|^2} < \frac{M}{\frac{d}{2} \cdot d^2} = \frac{2M}{d^3}$$

and, by the estimating theorem of contour integral,

$$|E| = |\Delta z| \cdot \left| \int_{\gamma} \frac{f(t)}{(t-z_1)(t-z)^2} dt \right| < |\Delta z| \cdot \frac{2M}{d^3} \cdot k,$$

where k is the length of the contour. The last expression tends to zero as $\Delta z \rightarrow 0$. This settles the proof.

Remark 1. By induction, one can prove the following generalisation of (2):

$$g^{(n)}(z) = n! \int_{\gamma} \frac{f(t)}{(t-z)^{n+1}} dt \quad (n = 0, 1, 2, \dots) \quad (3)$$

Remark 2. The contour γ may be . If it especially is a circle, then (1) defines a holomorphic function inside γ and another outside it.