



planetmath.org

Math for the people, by the people.

coefficients of Laurent series

Canonical name	CoefficientsOfLaurentSeries
Date of creation	2013-03-22 15:19:22
Last modified on	2013-03-22 15:19:22
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	15
Author	pahio (2872)
Entry type	Result
Classification	msc 30B10
Related topic	LaurentSeries
Related topic	TechniqueForComputingResidues
Related topic	UniquenessOfLaurentExpansion

Suppose that f is analytic in the annulus $\{z \in \mathbb{C} : R_1 < |z - a| < R_2\}$, where R_1 may be 0 and R_2 may be ∞ . Then the coefficients of the <http://planetmath.org/LaurentSeries> Laurent series

$$\sum_{n=-\infty}^{\infty} c_n (z - a)^n$$

of f can be obtained from

$$c_n = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(t)}{(t - a)^{n+1}} dt \quad (n = 0, \pm 1, \pm 2, \dots), \quad (1)$$

where the <http://planetmath.org/ContourIntegral> path γ goes anticlockwise once around the point $z = a$ within the annulus. Especially, the residue of f in the point a is

$$c_{-1} = \frac{1}{2\pi i} \oint_{\gamma} f(t) dt. \quad (2)$$

Remark. Usually, the Laurent series of a function, i.e. the coefficients c_n , are not determined by using the integral formula (1), but directly from known series. Often it is sufficient to know the value of c_{-1} or the residue, which is used to compute integrals (see the Cauchy residue theorem — cf. (2)). There is also the usable

Rule. In the case that the limit $\lim_{z \rightarrow a} (z - a)f(z)$ exists and has a non-zero value r , the point $z = a$ is a pole of the 1 for the function f and

$$\text{Res}(f; a) = r.$$

Examples

1. Let $f(z) := \frac{1}{\sin z}$, and $a = 0$. Using the Taylor series of the complex sine we obtain

$$\lim_{z \rightarrow 0} z \frac{1}{\sin z} = \lim_{z \rightarrow 0} \frac{1}{1 - \frac{z^2}{3!} + \dots} = 1,$$

whence $\text{Res}(\frac{1}{\sin z}; 0) = 1$. Thus we can write

$$\oint_{\gamma} \frac{dz}{\sin z} = 2\pi i,$$

where the γ must be chosen such that it encloses only the pole 0 of $\frac{1}{\sin z}$.

2. The Taylor series of the complex exponential function gives the Laurent series

$$e^{\frac{1}{z}} \equiv 1 + \frac{1}{z} + \frac{1}{2!z^2} + \frac{1}{3!z^3} + \dots$$

which shows that $\text{Res}(e^{\frac{1}{z}}; 0) = 1$.