



planetmath.org

Math for the people, by the people.

generalized Cauchy integral formula

Canonical name	GeneralizedCauchyIntegralFormula
Date of creation	2013-03-22 17:46:41
Last modified on	2013-03-22 17:46:41
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	6
Author	jirka (4157)
Entry type	Theorem
Classification	msc 30E20
Synonym	generalized Cauchy formula

**Theorem.** *Let  $U \subset \mathbb{C}$  be a domain with  $C^1$  boundary. Let  $f: U \rightarrow \mathbb{C}$  be a  $C^1$  function that is  $C^1$  up to the boundary. Then for  $z \in U$ ,*

$$f(z) = \frac{1}{2\pi i} \int_{\partial U} \frac{f(w)}{w - z} dw - \frac{1}{2\pi i} \int_U \frac{\frac{\partial f}{\partial \bar{z}}(w)}{w - z} d\bar{w} \wedge dw.$$

Note that  $C^1$  up to the boundary means that the function and the derivative extend to be continuous functions on the closure of  $U$ . The theorem follows from Stokes' theorem. When  $f$  is holomorphic, then the second term is zero and this is the classical Cauchy integral formula.

## References

- [1] Lars Hörmander. , North-Holland Publishing Company, New York, New York, 1973.
- [2] Steven G. Krantz. , AMS Chelsea Publishing, Providence, Rhode Island, 1992.