



planetmath.org

Math for the people, by the people.

evaluating the gamma function at $1/2$

Canonical name	EvaluatingTheGammaFunctionAt12
Date of creation	2013-03-22 16:57:13
Last modified on	2013-03-22 16:57:13
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	4
Author	CWoo (3771)
Entry type	Derivation
Classification	msc 30D30
Classification	msc 33B15
Related topic	AreaUnderGaussianCurve
Related topic	LaplaceTransformOfPowerFunction

In the entry on the gamma function it is mentioned that $\Gamma(1/2) = \sqrt{\pi}$. In this entry we reduce the proof of this claim to the problem of computing the area under the bell curve. First note that by definition of the gamma function,

$$\begin{aligned}\Gamma(1/2) &= \int_0^\infty e^{-x} x^{-1/2} dx \\ &= 2 \int_0^\infty e^{-x} \frac{1}{2\sqrt{x}} dx.\end{aligned}$$

Performing the substitution $u = \sqrt{x}$, we find that $du = \frac{1}{2\sqrt{x}} dx$, so

$$\Gamma(1/2) = 2 \int_0^\infty e^{-u^2} du = \int_{\mathbb{R}} e^{-u^2} du,$$

where the last equality holds because e^{-u^2} is an even function. Since the area under the bell curve is $\sqrt{\pi}$, it follows that $\Gamma(1/2) = \sqrt{\pi}$.