



Math for the people, by the people.

closure properties of Cauchy-Riemann equations

Canonical name	ClosurePropertiesOfCauchyRiemannEquations
Date of creation	2013-03-22 17:44:20
Last modified on	2013-03-22 17:44:20
Owner	rspuzio (6075)
Last modified by	rspuzio (6075)
Numerical id	14
Author	rspuzio (6075)
Entry type	Theorem
Classification	msc 30E99
Related topic	TangentialCauchyRiemannComplexOfCinfySmoothForms
Related topic	ACRcomplex

The set of solutions of the Cauchy-Riemann equations is closed under a surprisingly large number of operations. For convenience, let us introduce the notational conventions that f and g are complex functions with $f(x + iy) = u(x, y) + iv(x, y)$ and $g(x + iy) = p(x, y) + iq(x, y)$. Let D and D' denote open subsets of the complex plane.

Theorem 1. *If $f: D \rightarrow \mathbb{C}$ and $g: D \rightarrow \mathbb{C}$ satisfy the Cauchy-Riemann equations, so does $f + g$. Furthermore, if $z \in \mathbb{C}$, then zf satisfies the Cauchy-Riemann equations.*

Proof. This is an immediate consequence of the linearity of derivatives. \square

Theorem 2. *If $f: D \rightarrow \mathbb{C}$ and $g: D \rightarrow \mathbb{C}$ satisfy the Cauchy-Riemann equations, so does $f \cdot g$.*

Proof. Letting h and k denote the real and imaginary parts of $f \cdot g$ respectively, we have

$$\begin{aligned} \frac{\partial h}{\partial x} - \frac{\partial k}{\partial y} &= \frac{\partial}{\partial x} (up - vq) - \frac{\partial}{\partial y} (uq + vp) \\ &= u \frac{\partial p}{\partial x} + p \frac{\partial u}{\partial x} - v \frac{\partial q}{\partial x} - q \frac{\partial v}{\partial x} - u \frac{\partial q}{\partial y} - q \frac{\partial u}{\partial y} - v \frac{\partial p}{\partial y} - p \frac{\partial v}{\partial y} \\ &= u \left(\frac{\partial p}{\partial x} - \frac{\partial q}{\partial y} \right) - v \left(\frac{\partial p}{\partial y} + \frac{\partial q}{\partial x} \right) + p \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) - q \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right) = 0 \end{aligned}$$

and

$$\begin{aligned} \frac{\partial h}{\partial y} + \frac{\partial k}{\partial x} &= \frac{\partial}{\partial y} (up - vq) + \frac{\partial}{\partial x} (uq + vp) \\ &= u \frac{\partial p}{\partial y} + p \frac{\partial u}{\partial y} - v \frac{\partial q}{\partial y} - q \frac{\partial v}{\partial y} + u \frac{\partial q}{\partial x} + q \frac{\partial u}{\partial x} + v \frac{\partial p}{\partial x} + p \frac{\partial v}{\partial x} \\ &= u \left(\frac{\partial p}{\partial y} + \frac{\partial q}{\partial x} \right) + v \left(\frac{\partial p}{\partial x} - \frac{\partial q}{\partial y} \right) + p \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right) + q \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) = 0. \end{aligned}$$

\square

Theorem 3. *If $f: D \rightarrow D'$ and $g: D' \rightarrow \mathbb{C}$ satisfy the Cauchy-Riemann equations, so does $f \circ g$.*

Proof. Letting h and k denote the real and imaginary parts of $f \circ g$ respectively, we have

$$\begin{aligned}\frac{\partial h}{\partial x} - \frac{\partial k}{\partial y} &= \frac{\partial}{\partial x} u(p(x, y), q(x, y)) - \frac{\partial}{\partial y} v(p(x, y), q(x, y)) \\ &= \frac{\partial u}{\partial p} \frac{\partial p}{\partial x} + \frac{\partial u}{\partial q} \frac{\partial q}{\partial x} - \frac{\partial v}{\partial p} \frac{\partial p}{\partial y} - \frac{\partial v}{\partial q} \frac{\partial q}{\partial y} \\ &= \frac{\partial u}{\partial p} \left(\frac{\partial p}{\partial x} - \frac{\partial q}{\partial y} \right) + \frac{\partial q}{\partial y} \left(\frac{\partial u}{\partial p} - \frac{\partial v}{\partial q} \right) + \frac{\partial u}{\partial q} \left(\frac{\partial p}{\partial y} + \frac{\partial q}{\partial x} \right) - \frac{\partial p}{\partial y} \left(\frac{\partial u}{\partial q} + \frac{\partial v}{\partial p} \right) = 0\end{aligned}$$

and

$$\begin{aligned}\frac{\partial h}{\partial y} + \frac{\partial k}{\partial x} &= \frac{\partial}{\partial y} u(p(x, y), q(x, y)) + \frac{\partial}{\partial x} v(p(x, y), q(x, y)) \\ &= \frac{\partial u}{\partial p} \frac{\partial p}{\partial y} + \frac{\partial u}{\partial q} \frac{\partial q}{\partial y} + \frac{\partial v}{\partial p} \frac{\partial p}{\partial x} + \frac{\partial v}{\partial q} \frac{\partial q}{\partial x} \\ &= \frac{\partial u}{\partial p} \left(\frac{\partial p}{\partial y} + \frac{\partial q}{\partial x} \right) - \frac{\partial q}{\partial x} \left(\frac{\partial u}{\partial p} - \frac{\partial v}{\partial q} \right) - \frac{\partial u}{\partial q} \left(\frac{\partial p}{\partial x} - \frac{\partial q}{\partial y} \right) + \frac{\partial p}{\partial x} \left(\frac{\partial u}{\partial q} + \frac{\partial v}{\partial p} \right) = 0\end{aligned}$$

□

Theorem 4. *If $f: D \rightarrow \mathbb{C}$ satisfies the Cauchy-Riemann equations, and has non-vanishing Jacobian, then f^{-1} also satisfies the Cauchy-Riemann equations.*

Proof. Let us denote the real and imaginary parts of f^{-1} as h and k , respectively. Then, by definition of inverse function, we have

$$\begin{aligned}u(h(x, y), k(x, y)) &= x \\ v(h(x, y), k(x, y)) &= y.\end{aligned}$$

Taking derivatives,

$$\begin{aligned}\frac{\partial u}{\partial h} \frac{\partial h}{\partial x} + \frac{\partial u}{\partial k} \frac{\partial k}{\partial x} &= 1 \\ \frac{\partial u}{\partial h} \frac{\partial h}{\partial y} + \frac{\partial u}{\partial k} \frac{\partial k}{\partial y} &= 0 \\ \frac{\partial v}{\partial h} \frac{\partial h}{\partial x} + \frac{\partial v}{\partial k} \frac{\partial k}{\partial x} &= 0 \\ \frac{\partial v}{\partial h} \frac{\partial h}{\partial y} + \frac{\partial v}{\partial k} \frac{\partial k}{\partial y} &= 1\end{aligned}$$

By the Cauchy-Riemann equations, $\partial u/\partial h = \partial v/\partial k$ and $\partial u/\partial k = -\partial v/\partial h$. Using these relations to re-express the derivatives of u as derivatives of v , then subtracting the fourth equation from the first equation and adding the second and third equations, we obtain

$$\begin{aligned}\frac{\partial u}{\partial h} \left(\frac{\partial h}{\partial x} - \frac{\partial k}{\partial y} \right) + \frac{\partial u}{\partial k} \left(\frac{\partial h}{\partial y} + \frac{\partial k}{\partial x} \right) &= 0 \\ \frac{\partial u}{\partial h} \left(\frac{\partial h}{\partial y} + \frac{\partial k}{\partial x} \right) - \frac{\partial u}{\partial k} \left(\frac{\partial h}{\partial x} - \frac{\partial k}{\partial y} \right) &= 0.\end{aligned}$$

With a little algebraic manipulation, we may conclude

$$\begin{aligned}\left(\left(\frac{\partial u}{\partial h} \right)^2 + \left(\frac{\partial u}{\partial k} \right)^2 \right) \left(\frac{\partial h}{\partial y} + \frac{\partial k}{\partial x} \right) &= 0 \\ \left(\left(\frac{\partial u}{\partial h} \right)^2 + \left(\frac{\partial u}{\partial k} \right)^2 \right) \left(\frac{\partial h}{\partial x} - \frac{\partial k}{\partial y} \right) &= 0.\end{aligned}$$

Note that, by the Cauchy-Riemann equations, the Jacobian of f equals the common prefactor of these equations:

$$\frac{\partial(u, v)}{\partial(h, k)} = \frac{\partial u}{\partial h} \frac{\partial v}{\partial k} - \frac{\partial u}{\partial k} \frac{\partial v}{\partial h} = \left(\frac{\partial u}{\partial h} \right)^2 + \left(\frac{\partial u}{\partial k} \right)^2$$

Hence, by assumptions, this quantity differs from zero and we may cancel it to obtain the Cauchy-Riemann equations for f^{-1} . \square