



Math for the people, by the people.

Schwarz-Christoffel transformation

Canonical name	SchwarzChristoffelTransformation
Date of creation	2013-03-22 14:41:02
Last modified on	2013-03-22 14:41:02
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	15
Author	pahio (2872)
Entry type	Result
Classification	msc 31A99
Classification	msc 30C20
Synonym	Schwarz' transformation
Related topic	ConformalMapping

Let

$$w = f(z) = c \int \frac{dz}{(z - a_1)^{k_1} (z - a_2)^{k_2} \dots (z - a_n)^{k_n}} + C,$$

where the a_j 's are real numbers satisfying $a_1 < a_2 < \dots < a_n$, the k_j 's are real numbers satisfying $|k_j| \leq 1$; the integral expression means a complex antiderivative, c and C are complex constants.

The transformation $z \mapsto w$ maps the real axis and the upper half-plane <http://planetmath.org/ConformalMapping> conformally onto the closed area bounded by a broken line. Some vertices of this line may be in the infinity (the corresponding angles are $= 0$). When z moves on the real axis from $-\infty$ to ∞ , w moves along the broken line so that the direction turns the amount $k_j\pi$ anticlockwise every z passes a point a_j . If the broken line closes to a polygon, then $k_1 + k_2 + \dots + k_n = 2$.

This transformation is used in solving two-dimensional potential problems. The parameters a_j and k_j are chosen such that the given polygonal domain in the complex w -plane can be obtained.

A half-trivial example of the transformation is

$$w = \frac{1}{2} \int \frac{dz}{(z - 0)^{\frac{1}{2}}} = \sqrt{z},$$

which maps the upper half-plane onto the first quadrant of the complex plane.