



Math for the people, by the people.

## CR function

Canonical name	CRFunction
Date of creation	2013-03-22 14:57:10
Last modified on	2013-03-22 14:57:10
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	5
Author	jirka (4157)
Entry type	Definition
Classification	msc 32V10
Defines	CR distribution

**Definition.** Let  $M \subset \mathbb{C}^N$  be a CR submanifold and let  $f$  be a  $C^k(M)$  ( $k$  times continuously differentiable) function to  $\mathbb{C}$  where  $k \geq 1$ . Then  $f$  is a *CR function* if for every CR vector field  $L$  on  $M$  we have  $Lf \equiv 0$ . A <http://planetmath.org/Distribution4distribution>  $f$  on  $M$  is called a *CR distribution* if similarly every CR vector field annihilates  $f$ .

For example restrictions of holomorphic functions in  $\mathbb{C}^N$  to  $M$  are CR functions. The converse is not always true and is not easy to see. For example the following basic theorem is very useful when you have real analytic submanifolds.

**Theorem.** *Let  $M \subset \mathbb{C}^N$  be a generic submanifold which is real analytic (the defining function is real analytic). And let  $f: M \rightarrow \mathbb{C}$  be a real analytic function. Then  $f$  is a CR function if and only if  $f$  is a restriction to  $M$  of a holomorphic function defined in an open neighbourhood of  $M$  in  $\mathbb{C}^N$ .*

## References

- [1] M. Salah Baouendi, Peter Ebenfelt, Linda Preiss Rothschild. , Princeton University Press, Princeton, New Jersey, 1999.